

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
22 January 2004 (22.01.2004)

PCT

(10) International Publication Number  
WO 2004/007670 A2

(51) International Patent Classification<sup>7</sup>: C12N

(21) International Application Number:  
PCT/US2003/021570

(22) International Filing Date: 10 July 2003 (10.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/395,431 12 July 2002 (12.07.2002) US

(71) Applicant: NEW ENGLAND BIOLABS, INC.  
[US/US]; 32 Tozer Road, Beverly, MA 01915 (US).

(72) Inventors: MORGAN, Richard, D.; 31 Donovan's  
Way, Middleton, MA 01949 (US). BHATIA, Tanya;  
504 Pinebrook Drive, Peabody, MA 01960 (US). DAVIS,  
Theodore; 139 Depot Road, Boxford, MA 01921 (US).  
LOVASCO, Lindsay; 4 Sutton Avenue, Salem, MA  
01970 (US).

(74) Common Representative: WILLIAMS, Gregory, D. et  
al.; New England Biolabs, Inc., 32 Tozer Road, Beverly,  
MA 01915 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— as to the applicant's entitlement to claim the priority of the  
earlier application (Rule 4.17(iii)) for all designations

**Published:**

— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: RECOMBINANT TYPE II RESTRICTION ENDONUCLEASES, MmeI AND RELATED ENDONUCLEASES AND  
METHODS FOR PRODUCING THE SAME

(57) Abstract: In accordance with the present invention, there is provided a DNA (deoxyribonucleic acid) fragment which encodes  
the MmeI type II restriction endonuclease enzyme. This one polypeptide possesses two related enzymatic functions; namely an  
endonuclease activity which recognizes the DNA sequence 5'-TCC(Pu)AC-3' and cleaves as indicated by the arrows: 5' -TCCRAC  
(N20) ↓-3' 3'-AGGYTG(N18)↑-5' and a second enzymatic activity that recognizes the same DNA sequence, 5'-TCC(Pu)AC-3',  
but modifies this sequence by the addition of a methyl group to prevent cleavage by the MmeI endonuclease activity.

WO 2004/007670 A2

**RECOMBINANT TYPE II RESTRICTION ENDONUCLEASES,  
MmeI AND RELATED ENDONUCLEASES AND METHODS FOR PRODUCING  
THE SAME**

5

**BACKGROUND OF THE INVENTION**

The present invention relates to a DNA (deoxyribonucleic acid) fragment, which fragment codes for one polypeptide possessing two related enzymatic functions, namely an enzyme which recognizes the DNA sequence 5'-TCC(Pu)AC-3' and cleaves the phosphodiester bond between the 20th and 21st residues 3' to this recognition sequence on this DNA strand, and between the 18th and 19th residues 5' to the recognition sequence on the complement strand 5'-GT(Py)GGT-3' to produce a 2 base 3' extension (hereinafter referred to as the MmeI restriction endonuclease), and a second enzymatic activity that recognizes the same DNA sequence, 5'-TCC(Pu)AC-3', but modifies this sequence by the addition of a methyl group to prevent cleavage by the MmeI endonuclease. The present invention also relates to a vector containing the DNA fragment, a transformed host containing this DNA fragment, and an improved process for producing MmeI restriction endonuclease from such a transformed host. The present invention also relates to a process for identifying additional DNA fragments that encode enzymes having the same general properties as MmeI but potentially having unique DNA recognition sequences. This process depends on the use of the amino acid sequence of the MmeI enzyme presented in this application, or subsequently on the additional sequences identified through this process. The invention also relates to additional DNA fragments, identifiable through the process described, each of which encodes a polypeptide having significant amino acid sequence similarity to the MmeI polypeptide. The polypeptides

10

15

20

25

30

35

-2-

5 encoded by these DNA fragments are predicted to perform  
similar functions to MmEI. Specifically, they are  
predicted to possess the dual enzymatic functions of  
cleaving DNA in a specific manner at a relatively far  
distance from the specific recognition sequence and also  
modifying their recognition sequences to protect the  
host DNA from cleavage by endonuclease activity. An  
example of such an enzyme identified by this process is  
CstMI (see U.S. Application Serial No.:\_\_\_\_\_, filed  
10 concurrently herewith). CstMI was identified as a  
potential endonuclease because of its highly significant  
amino acid sequence similarity to MmEI. CstMI recognizes  
the sequence 5'-AAGGAG-3' and cleaves the phosphodiester  
bond between the 20th and 21st residues 3' to the  
15 recognition sequence on this DNA strand, and between the  
18th and 19th residues 5' to the recognition sequence on  
the complement strand 5'-CTCCTT-3' to produce a 2 base  
3' extension.

20 Restriction endonucleases are a class of enzymes  
that occur naturally in prokaryotes. There are several  
classes of restriction systems known, of which the type  
II endonucleases are the class useful in genetic  
engineering. When these type II endonucleases are  
25 purified away from other contaminating prokaryotic  
components, they can be used in the laboratory to break  
DNA molecules into precise fragments. This property  
enables DNA molecules to be uniquely identified and to  
be fractionated into their constituent genes.  
30 Restriction endonucleases have proved to be  
indispensable tools in modern genetic research. They  
are the biochemical 'scissors' by means of which genetic  
engineering and analysis is performed.

-3-

Restriction endonucleases act by recognizing and binding to particular sequences of nucleotides (the 'recognition sequence') along the DNA molecule. Once bound, the type II endonucleases cleave the molecule within, or to one side of, the sequence. Different restriction endonucleases have affinity for different recognition sequences. The majority of restriction endonucleases recognize sequences of 4 to 6 nucleotides in length, although recently a small number of restriction endonucleases which recognize 7 or 8 uniquely specified nucleotides have been isolated. Most recognition sequences contain a dyad axis of symmetry and in most cases all the nucleotides are uniquely specified. However, some restriction endonucleases have degenerate or relaxed specificities in that they recognize multiple bases at one or more positions in their recognition sequence, and some restriction endonucleases recognize asymmetric sequences. *HaeIII*, which recognizes the sequence 5'-GGCC-3', is an example of a restriction endonuclease having a symmetrical, non-degenerate recognition sequence; *HaeII*, which recognizes 5'-(Pu)GCGC(Py)-3' typifies restriction endonucleases having a degenerate or relaxed recognition sequence; while *BspMI*, which recognizes 5'-ACCTGC-3' typifies restriction endonucleases having an asymmetric recognition sequence. Type II endonucleases with symmetrical recognition sequences generally cleave symmetrically within or adjacent to the recognition site, while those that recognize asymmetric sequences tend to cleave at a distance of from 1 to 20 nucleotides to one side of the recognition site. The enzyme of this application, *MmeI*, (along with *CstMI*) has the distinction of cleaving the DNA at the farthest distance from the recognition sequence of any known type II restriction endonuclease. More than two hundred unique

-4-

restriction endonucleases have been identified among several thousands of bacterial species that have been examined to date.

5           A second component of restriction systems are the modification methylases. These enzymes are complementary to restriction endonucleases and they provide the means by which bacteria are able to protect their own DNA and distinguish it from foreign, infecting  
10       DNA. Modification methylases recognize and bind to the same nucleotide recognition sequence as the corresponding restriction endonuclease, but instead of breaking the DNA, they chemically modify one or other of the nucleotides within the sequence by the addition of a  
15       methyl group. Following methylation, the recognition sequence is no longer cleaved by the restriction endonuclease. The DNA of a bacterial cell is modified by virtue of the activity of its modification methylase and it is therefore insensitive to the presence of the  
20       endogenous restriction endonuclease. It is only unmodified, and therefore identifiably foreign, DNA that is sensitive to restriction endonuclease recognition and cleavage. Modification methyltransferases are usually separate enzymes from their cognate endonuclease  
25       partners. In some cases, there is a single polypeptide that possesses both a modification methyltransferase function and an endonuclease function, for example, Eco57I. In such cases, there is a second methyltransferase present as part of the restriction-  
30       modification system. In contrast, the MmeI system of the present application has no second methyltransferase accompanying the endonuclease-methyltransferase polypeptide.

-5-

Endonucleases are named according to the bacteria from which they are derived. Thus, the species *Haemophilus aegyptius*, for example synthesizes 3 different restriction endonucleases, named *HaeI*, *HaeII* and *HaeIII*. These enzymes recognize and cleave the sequences 5'-(W)GGCC(W)-3', 5'-(Pu)GCGC(Py)-3' and 5'-GGCC-3' respectively. *Escherichia coli* RY13, on the other hand, synthesizes only one enzyme, *EcoRI*, which recognizes the sequence 5'-GAATTC-3'.

While not wishing to be bound by theory, it is thought that in nature, restriction endonucleases play a protective role in the welfare of the bacterial cell. They enable bacteria to resist infection by foreign DNA molecules such as viruses and plasmids that would otherwise destroy or parasitize them. They impart resistance by binding to infecting DNA molecules and cleaving them in each place that the recognition sequence occurs. The disintegration that results inactivates many of the infecting genes and renders the DNA susceptible to further degradation by exonucleases.

More than 3000 restriction endonucleases have been isolated from various bacterial strains. Of these, more than 240 recognize unique sequences, while the rest share common recognition specificities. Restriction endonucleases which recognize the same nucleotide sequence are termed "isoschizomers." Although the recognition sequences of isoschizomers are the same, they may vary with respect to site of cleavage (e.g., *XmaI* v. *SmaI*, Endow, et al., *J. Mol. Biol.* **112**:521 (1977); Waalwijk, et al., *Nucleic Acids Res.* **5**:3231 (1978)) and in cleavage rate at various sites (*XhoI* v. *PaeR7I*, Gingeras, et al., *Proc. Natl. Acad. Sci. U.S.A.* **80**:402 (1983)).

-6-

Restriction endonucleases have traditionally been classified into three major classes; type I, type II and type III. The type I restriction systems assemble a multi-peptide complex consisting of restriction polypeptide, modification polypeptide, and specificity, or DNA recognition, polypeptide. Type I systems require a divalent cation, ATP and S-adenylosyl-methionine (SAM) as cofactors. Type I systems cleave DNA at random locations up to several thousand basepairs away from their specific recognition site. The type III systems generally recognize an asymmetric DNA sequence and cleave at a specific position 20 to 30 basepairs to one side of the recognition sequence. Such systems require the cofactor ATP in addition to SAM and a divalent cation. The type III systems assemble a complex of endonuclease polypeptide and modification polypeptide that either modifies the DNA at the recognition sequence or cleaves. Type III systems produce partial digestion of the DNA substrate due to this competition between their modification and cleavage activities, and so have not been useful for genetic manipulation.

MmeI does not require ATP for DNA cleavage activity and it cleaves to completion; thus it can be classified as a type II endonuclease. Unlike other type II enzymes, however, MmeI consists of a single polypeptide that combines both endonuclease and modification activities and is sufficient by itself to form the entire restriction modification system. MmeI also cleaves the farthest distance from the specific DNA recognition sequence of any type II endonuclease (as does CstMI of this application). MmeI is quite large and appears to have three functional domains combined in one polypeptide. These consist of an amino-terminal domain which contains the endonuclease DNA cleavage motif and



-7-

which may also be involved in DNA recognition, a DNA modification domain most similar to the gamma-class N6mA methyltransferases, and a carboxy-terminal domain presumed to be involved in dimer formation and possibly DNA recognition. The enzyme requires SAM for both cleavage and modification activity. The single MmeI polypeptide is sufficient to modify the plasmid vector carrying the gene *in vivo* to provide protection against MmeI cleavage *in vitro*, yet it is also able to cleave unmodified DNAs *in vitro* when using the endonuclease buffer containing Mg++ and SAM.

There is a continuing need for novel type II restriction endonucleases. Although type II restriction endonucleases which recognize a number of specific nucleotide sequences are currently available, new restriction endonucleases which recognize novel sequences provide greater opportunities and ability for genetic manipulation. Each new unique endonuclease enables scientists to precisely cleave DNA at new positions within the DNA molecule, with all the opportunities this offers.

#### SUMMARY OF THE INVENTION

In accordance with the present invention, there is provided a novel DNA fragment encoding a novel restriction endonuclease, obtainable from *Methylophilus methylotrophus* (NEB#1190). The endonuclease is hereinafter referred to as "MmeI", which endonuclease:

- (1) recognizes the degenerate nucleotide sequence 5'-TCC(Pu)AC-3' in a double-stranded DNA molecule as shown below:

-8-

5'-TCC(Pu)AC-3'

3'-AGG(Py)TG-5'

(wherein G represents guanine, C represents cytosine, A represents adenine, T represents thymine, (Pu) represents a purine, either A or G, and (Py) represents a pyrimidine, either C or T);

- 5
- 10 (2) cleaves DNA in the phosphodiester bond following the 20th nucleotide 3' to the recognition sequence 5'-TCC(Pu)AC-3 and preceding the 18th nucleotide 5' to the complement strand of the recognition sequence
- 15 5'-GT(Py)GGA-3' to produce a 2 base 3' extension:

5'-TCC(Pu)AC(N20)/-3'

3'-AGG(Py)GT(N18)/-5'; and

- 20
- (3) methylates the recognition sequence specified in (1) *in vivo* to protect the host DNA from cleavage by the MmeI endonuclease activity;

25 The invention further relates to additional DNA fragments, each of which is identified to encode polypeptides which share significant sequence similarity to the MmeI restriction-modification polypeptide. The DNA fragment encoding the MmeI polypeptide enables the

30 identification of these additional potential endonucleases by using similarity searching of the MmeI sequence against sequences available in databases, such as GENBANK, using a program such as BLAST (Altschul, et al. Nucleic Acids Res. 25:3389-3402 (1997)). These DNA

35 fragments, as well as any other fragments with such

-9-

similarity to MmeI that may be deposited in the databases in the future, are candidates which may encode polypeptides that are similar to MmeI, in that the polypeptides encoded act as both restriction  
5 endonuclease and methyltransferase. These polypeptides may, like MmeI, cleave DNA at a similarly far distance from the recognition sequence, in the range of 18 to 20 nucleotides or more, which character is unique and useful in certain molecular biology technologies.  
10 Specifically these polypeptides contain amino acid motifs common to N6mA DNA methyltransferases in the middle of the polypeptide, have a motif common to restriction endonucleases and located in the amino-terminal section of the polypeptides, consisting of the  
15 amino acids D/E(X8-X12)D/EXK, and have a region of several hundred amino acids following the conserved methyltransferase motifs which are significantly similar to this region of MmeI and are believed to serve as a dimerization and possibly a DNA sequence recognition  
20 domain. An example of such a polypeptide, CstMI, is presented. CstMI has been shown to recognize the 6 base pair asymmetric sequence 5'-AAGGAG-3' and to cleave the DNA in the same manner as MmeI; 5'-AAGGAGN20/N18-3'. The endonuclease encoded by these DNA fragments may be  
25 produced by the process used for MmeI, as described below.

The present invention further relates to a process for the production of the restriction endonuclease MmeI.  
30 This process comprises culturing a transformed host, such as *E. coli*, containing the DNA fragment encoding the MmeI restriction system polypeptide, collecting the cultured cells, obtaining a cell-free extract therefrom and separating and collecting the restriction  
35 endonuclease MmeI from the cell-free extract. The

-10-

present invention further relates to a process for the production of the restriction endonucleases encoded by the DNA sequences identified as homologous to MmeI. This process comprises culturing a transformed host, such as *E. coli*, containing the gene for these restriction systems, collecting the cultured cells, obtaining a cell-free extract therefrom and separating and collecting the restriction endonuclease from the cell-free extract.

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 - Agarose gel showing MmeI cleavage of lambda, T7, phiX174, pBR322 and pUC19 DNAs.

Figure 2 - DNA sequence of the MmeI gene locus (SEQ ID NO:1).

Figure 3 - Amino acid sequence of the MmeI gene locus (SEQ ID NO:2).

Figure 4 - Agarose gel showing MmeI cleavage of pTBMmeI.1 DNA and unmodified DNA substrates.

Figure 5 - Agarose gel showing MmeI cleavage of unmethylated, hemi-methylated and fully methylated DNA substrates.

Figure 6 - Incorporation of labeled methyl groups into unmethylated, hemi-methylated and fully methylated DNA substrates.

Figure 7 - Multiple sequence alignment of MmeI amino acid sequence (SEQ ID NO:3 through SEQ ID NO:14) and homologous polypeptides from public databases.

-11-

DETAILED DESCRIPTION OF THE INVENTION

5       The recognition sequence and cleavage site of the  
endoneuclease of the present invention were previously  
described (Boyd, Nucleic Acids Res. 14: 5255-5274  
(1986)). However the MmeI enzyme proved difficult to  
produce from the native host, *Methylophilus*  
10 *methylophilus*, due to very low yield of the enzyme and  
the relative difficulty of growing the *M. methylophilus*  
host in large quantity. To overcome these limitations to  
producing MmeI, the present application describes the  
identification of the DNA sequence encoding the MmeI  
15 gene and the expression of this MmeI gene in a suitable  
host, in the present instance *E. coli*. This manipulation  
of the MmeI encoding DNA fragment results in both a  
significant increase in the amount of enzyme produced  
per gram of cells and a significant increase in ease of  
growth of large amounts of cells containing MmeI enzyme.

20       Several standard approaches typically employed by  
persons skilled in the art of cloning were applied to  
the task of cloning of MmeI without success.  
Specifically, the methylase selection approach (Wilson,  
25 et al., U.S. Patent No. 5,200,333) was attempted  
unsuccessfully. Several random libraries of *M.*  
*methylophilus* DNA were constructed in *E. coli* and  
challenged by digesting with MmeI, but no MmeI methylase  
containing clones were obtained.

30       A second approach was also attempted but failed. In  
this approach, antibodies specific for N6mA were used to  
screen a library of random clones constructed in a  
lambda phage replacement vector. The approach was  
35 successful in obtaining methylase positive clones, but

-12-

all examined were found to express the methyltransferase of the second restriction system in *M. methylotrophus*, the MmeII methylase (recognition sequence 5'-GATC-3') rather than the desired MmeI methylase activity.

5

The successful approach to obtain the desired DNA fragment encoding the MmeI restriction system involved several steps. First a novel purification procedure was developed to purify the MmeI endonuclease peptide to  
10 homogeneity from *M. methylotrophus*. Once this ultra pure MmeI endonuclease polypeptide was successfully obtained in a significant amount, amino acid sequence from the amino terminus and from internal cyanogen-bromide  
15 degradation peptides was determined. Using the amino acid sequence obtained, degenerate DNA primers complementary to the DNA coding for the amino acid sequences were synthesized and used to PCR amplify a portion of the MmeI gene. The DNA sequence of this  
20 portion of the MmeI gene was determined. The entire MmeI endonuclease gene and surrounding DNA sequences were then obtained by applying the inverse PCR technique. A number of primers matching the DNA sequence obtained were designed, synthesized and used in combination with  
25 numerous different templates. The inverse PCR templates were produced by digesting *M. methylotrophus* genomic DNA with various restriction endonucleases and then ligating the cut *M. methylotrophus* DNA at low concentration to obtain circular molecules. The various primers were  
30 tried in combinations with the various templates to find primer-template combinations that produced a specific PCR amplification product. The products thus obtained were sequenced. Once the DNA sequence encoding the entire MmeI endonuclease gene was obtained, primers were  
35 designed to specifically amplify the gene from *M. methylotrophus* genomic DNA. The amplified gene was

-13-

inserted into an expression vector and cloned into an *E. coli* host. The host was tested and found to both express MmeI endonuclease activity and to *in vivo* modify the recombinant expression vector such that it was protected against MmeI endonuclease activity *in vitro*.

This finding that the single polypeptide encoding the MmeI endonuclease also provided *in vivo* protection against MmeI is in contrast to the previously published information on MmeI (Tucholski, Gene 223:293-302 (1998)). Specifically, this reference taught that the MmeI endonuclease polypeptide did not provide protection against MmeI endonuclease cleavage. This reference reported a separate methyltransferase of 48kD as required to modify the MmeI site on both strands and thus block cleavage by the MmeI endonuclease. Specifically, the reference teaches that the MmeI endonuclease polypeptide modifies the adenine in the top strand of the recognition sequence only, 5'-TCCRAC-3' and that such modified DNA is cut by the MmeI endonuclease. The DNA fragment of the present invention encodes the MmeI endonuclease gene, which when grown alone in an *E. coli* host renders the vector containing the MmeI endonuclease resistant to cleavage by the purified MmeI endonuclease. Further, the MmeI endonuclease produced from this fragment does not cleave a DNA fragment modified at the adenine of the top strand, 5'-TCCRAC-3' when no modification of the opposite, or bottom strand is present. This is in contrast to the teaching of the Tucholski reference. Also, the MmeI endonuclease of this application does cleave a DNA fragment in which the adenine residue in the bottom strand is modified 5'-GTYGGA-3' in contrast to the teaching of the Tucholski reference. When both the top strand and the bottom strand are modified at the

-14-

adenine residues, the MmeI endonuclease does not cleave the DNA. No second methyltransferase gene, such as reported in the Tucholski reference, was found adjacent to the MmeI endonuclease gene. There is an open reading frame immediately 3' to the MmeI endonuclease gene which would encode a protein of approximately the reported size of such a second methyltransferase activity (48kD). However, this potential polypeptide does not have the amino acid motifs found in methyltransferases, nor did it provide protection against MmeI endonuclease when cloned in *E. coli*. While the Tulchoski reference taught the necessity of a second methyltransferase polypeptide to provide protection against MmeI endonuclease activity for the host cell, it is demonstrated in the present application that the DNA fragment encoding the MmeI endonuclease polypeptide is sufficient to provide such protection. Additionally, the eleven DNA fragments described herein which encode amino acid sequences similar to MmeI are not flanked by any recognizable DNA methyltransferase genes. This indicates that these polypeptides are also likely to provide both protection for the host DNA and endonuclease activity against unmodified DNA substrates on their own, without having a second methyltransferase as part of the restriction modification system. This contrasts with other type II restriction modification systems.

The same group (Tucholski, Gene 223: 293-302 (1998), and Anna Podhajska, personal communication) had previously reported an amino acid sequence of eight residues for a single internal CnBr digestion fragment (sequence GRGRGVGV (SEQ ID NO:\_\_\_\_)). PCR based on this sequence was attempted yet failed repeatedly. This sequence was found to be unrelated to MmeI once the actual MmeI amino acid sequence was determined in



-15-

accordance with the present invention. Therefore correct internal amino acid sequences determination, which enabled the cloning of the MmEI gene, depended on the novel purification method described in this application for the production of sufficiently pure MmEI in large enough quantity to determine cyanogen bromide internal fragment amino acid sequences, as performed in this Application.

In Example II we obtained MmEI by culturing a transformed host carrying the MmEI gene, such as *E. coli* ER2683 carrying pTBMmEI.1 and recovering the endonuclease from the cells. A sample of *E. coli* ER2683 carrying pTBMmEI.1 (NEB#1457) has been deposited under the terms and conditions of the Budapest Treaty with the American Type Culture Collection (ATCC) on July 3, 2002 and bears the Patent Accession No. PTA-4521.

For recovering the enzyme of the present invention *E. coli* carrying pTBMmEI.1 (NEB#1457) may be grown using any suitable technique. For example, *E. coli* carrying pTBMmEI.1 may be grown in Luria broth media containing 100µg/ml ampicillin and incubated aerobically at 37°C with aeration. Cells in the late logarithmic stage of growth are induced by adding 0.3mM IPTG, grown for an additional 4 hours, collected by centrifugation and either disrupted immediately or stored frozen at -70°C.

The MmEI enzyme can be isolated from *E. coli* carrying pTBMmEI.1 cells by conventional protein purification techniques. For example, cell paste is suspended in a buffer solution and treated by sonication, high pressure dispersion or enzymatic digestion to allow extraction of the endonuclease by the buffer solution. Intact cells and cellular debris are

-16-

then removed by centrifugation to produce a cell-free extract containing MmeI. The MmeI endonuclease, along with its corresponding intrinsic methylase activity, is then purified from the cell-free extract by ion-exchange chromatography, affinity chromatography, molecular sieve chromatography, or a combination of these methods to produce the endonuclease of the present invention.

The present invention also relates to methods for identifying additional DNA fragments, each of which encodes a polypeptide having significant amino acid sequence similarity to the MmeI polypeptide. The polypeptides encoded by these DNA fragments are predicted to perform similar functions to MmeI. Specifically, they are predicted to possess the dual enzymatic functions of cleaving DNA in a specific manner at a relatively far distance from the specific recognition sequence and also modifying their recognition sequences to protect the host DNA from cleavage by their endonuclease activity. Once the amino acid sequence of the MmeI endonuclease was determined as described in this application, sequences deposited in databases can be compared to this MmeI sequence to find those few sequences that are highly significantly similar to MmeI. This method is similar to that of U.S. Patent No. 6,383,770 (Roberts, et al.), except that here we are searching for similarity to the MmeI endonuclease sequence, rather than searching for sequences that match a database of methyltransferase or endonuclease proteins and then examining any unidentified open reading frames next to potential methyltransferase open reading frames. Prior to identifying the MmeI amino acid sequence, the DNA sequences coding for proteins related to MmeI had not been included in the database of restriction and methyltransferase gene sequences utilized by Roberts, et al., *supra* since these sequences had not been linked to

-17-

any known endonuclease function. The method disclosed herein of identifying potential MmeI-like endonucleases is thus more specific than the method of U.S. Patent No. 6,383,770 (Roberts, et al.).

5

Similarity searching of the MmeI sequence against sequences available in databases, such as GENBANK, is accomplished using a program such as BLAST (Altschul, et al. Nucleic Acids Res. 25:3389-3402 (1997)). A sequence with an expectation value (E) score of less than  $E = e^{-10}$  is considered a potential candidate endonuclease.

10

Sequences that give expectation values that are much lower, such as less than  $E = e^{-30}$  is to be considered as highly likely to be endonucleases like MmeI. Such candidate MmeI-like peptides are further examined to see if they conform to the domain architecture that MmeI exhibits. A true candidate will contain an endonuclease fold motif, usually of the form (D/E)X8-X12(D/E)XK in the amino-terminal portion of the peptide, (Aravind et al. Nucleic Acid Res. 28:3417-3432 (2000)). A true candidate will contain methyltransferase motifs in the middle portion of the peptide similar to gamma class N6-methyl adenine methyltransferases, and sequences similar to the carboxyl portion of MmeI in the carboxyl portion of the candidate peptide. Such a BLAST search performed on June 12, 2003 returned the following sequences as highly significantly similar to MmeI:

15

20

25

Docket No.: NEB-207-PCT

Genbank accession ID      Description      Score      E value

GENBANK ACCESSION NO.	DESCRIPTION	SCORE	E VALUE	SEQ ID NO:
1. gi 15794682 ref NP_284504.1	hypothetical protein [Neisseri	643	0.0	6
2. gi 9945797 gb AAG03371.1	GcrY [Corynebacterium striatum	604	e-171	8
3. gi 16077744 ref NP_388558.1	similar to hypothetical protei	564	e-159	7
4. gi 28373198 ref NP_783835.1	putative YeeA protein [Lactoba...	531	e-149	3
5. gi 23110638 gb ZP_00096791.1	hypothetical protein [Novosph...	426	e-118	10
6. gi 27450519 gb AAO14619.1 AF465251_62	unknown [Lactobacillus...	217	9e-55	4
7. gi 15807258 ref NP_295988.1	DNA modification methyltransfe...	213	1e-53	14
8. gi 15807788 ref NP_285443.1	conserved hypothetical protein...	164	7e-39	13
9. gi 21231551 ref NP_637468.1	conserved hypothetical protein...	142	2e-32	N/A
10. gi 20803963 emb CAD31540.1	PUTATIVE DNA METHYLASE PROTEIN ...	134	7e-30	11
11. gi 23451826 gb AAN32874.1 AF461726_1	unknown [Pseudomonas f...	98	6e-19	9
12. gi 16125079 ref NP_419643.1	conserved hypothetical protein...	92	3e-17	12
13. gi 10954534 ref NP_044172.1	M. jannaschii predicted coding...	76	2e-12	N/A

-19-

Most of these proteins are labeled as hypothetical or putative in their database entries. A number of these appear to be full-length polypeptides, such as sequence #2 above: GcrY. Such candidates can be expressed as described in Roberts to identify the expected endonuclease activity. Some endonuclease genes may be inactive in the particular strain used for sequencing (Lin, et al. Proc. Natl. Acad. Sci. USA 98:2740-2745 (2001)). In such a circumstance it may prove possible to express functional endonucleases by repairing the mutations that have inactivated these genes. Several of the MmeI homologs, such as #7 (SEQ ID NO:14) (Deinococcus radiodurans DR2267) and #8 (SEQ ID NO:13) (Deinococcus radiodurans DR0119.1) have disruptions in the open reading frames. DR2267 has a stop codon, TAG, which prematurely terminates the open reading frame, in a position where MmeI has a glutamate amino acid coded for by the codon GAG. By changing this TAG stop codon to GAG it may be possible to reactivate this potential endonuclease gene. DR0119.1 is also disrupted, in that it has a frameshift that disrupts open reading frame. The MmeI sequence may be used as a guide to direct where to repair this frameshift by maximizing the similarity of the DR0119.1 sequence to the MmeI sequence. This may well restore DR0119.1 endonuclease activity.

An alternative way to generate potential new endonucleases is to take advantage of their similar domain structure by performing domain swapping. One may be able to swap the amino terminal domain of an MmeI-like peptide, for the amino terminal domain in the MmeI protein, for example by swapping the sequence of the potential new gene up to the first methyltransferase motif (motif X, "Gly Ala His Tyr Thr Ser" into MmeI to replace this portion of MmeI up to the same sequence.

-20-

This approach may be particularly useful when only a partial sequence is available or a potential gene has lost function due to multiple mutations. This approach will create a chimeric protein that potentially has endonuclease activity and cleaves at a distance away from the recognition sequence, like MmeI, but that recognizes a novel DNA sequence. One may also find sequences in the databases that are highly similar to MmeI but that are partial. For example, sequence #11 (SEQ ID NO:9) above (*Pseudomonas fluorescens*) is from a small fragment of DNA sequence in the database. To obtain a functional endonuclease like MmeI from this sequence one can use inverse PCR or other techniques to obtain DNA sequence adjacent to the fragment reported, then use that sequence to obtain an intact endonuclease gene.

Once a sequence is identified the potential endonuclease can be expressed and characterized as described in Roberts, et al. *supra*. Here, however, there is no separate methyltransferase gene to express along with the endonuclease. Once such a potential endonuclease is cloned and expressed in a suitable host, such as in *E. coli*, a cell free extract is prepared and analyzed to detect any endonuclease activity. Such an endonuclease assay must include the SAM cofactor required by these endonucleases. Once specific DNA cleavage activity is found the recognition sequence and cleavage site may be determined by standard methods. (Schildkraut, (1984) In Genet. Eng. (N Y) Vol 6. (Setlow J.K., Hollaender, A. Ed.). pp 117-140. Plenum Press, New York. "Screening for and characterizing restriction endonucleases.")

-21-

The enzymes so identified can be isolated from *E. coli* cells carrying the DNA fragment in a suitable vector by conventional protein purification techniques. For example, cell paste is suspended in a buffer solution and treated by sonication, high pressure dispersion or enzymatic digestion to allow extraction of the endonuclease by the buffer solution. Intact cells and cellular debris are then removed by centrifugation to produce a cell-free extract containing the enzyme. The endonuclease, along with its corresponding intrinsic methylase activity, is then purified from the cell-free extract by ion-exchange chromatography, affinity chromatography, molecular sieve chromatography, or a combination of these methods to produce the endonuclease of the present invention.

These DNA fragments, as well as any other fragments with such similarity to *MmeI* that may be deposited in the databases in the future, are predicted to encode polypeptides that are similar to *MmeI*, in that the polypeptides encoded act as both restriction endonuclease and methyltransferase. These polypeptides may, like *MmeI*, cleave DNA at a similarly far distance from the recognition sequence, in the range of about 18 to 20 nucleotides or more, which character is unique and useful in certain molecular biology technologies.

An example of such an enzyme identified by this process is *CstMI* (see U.S. Application Serial No. \_\_\_\_\_, filed concurrently herewith). *CstMI* was identified as a potential endonuclease because of its highly significant amino acid sequence similarity to *MmeI*. *CstMI* is encoded by sequence #2 above (SEQ ID NO:8), which gave highly significant Expectation value of  $e^{-171}$  when compared to *MmeI* by BLAST. *CstMI* recognizes

-22-

the 6 base pair asymmetric sequence 5'-AAGGAG-3' and cleaves the DNA in the same manner as MmeI: it cleaves the phosphodiester bond between the 20th and 21st residues 3' to this recognition sequence on this DNA strand, and between the 18th and 19th residues 5' to the recognition sequence on the complement strand 5'-CTCCTT-3' to produce a 2 base 3' extension.

The present invention is further illustrated by the following Examples. These Examples are provided to aid in the understanding of the invention and are not construed as a limitation thereof.

The references cited above and below are herein incorporated by reference.

#### EXAMPLE I

##### **PURIFICATION OF MmeI ENDONUCLEASE**

A single colony of *Methylophilus methylotrophus* (NEB#1190) was grown for 24 hrs in 1 liter of medium M (0.08  $\mu$ M CuSO<sub>4</sub>, 0.448  $\mu$ M MnSO<sub>4</sub>, 0.348  $\mu$ M ZnSO<sub>4</sub>, 6.0  $\mu$ M FeCl<sub>3</sub>, 18  $\mu$ M CaCO<sub>3</sub>, 1.6 mM MgSO<sub>4</sub>, 9.0 mM NaH<sub>2</sub>PO<sub>4</sub>, 10.9 mM K<sub>2</sub>HPO<sub>4</sub>, 13.6 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>) for 24 hours. This culture was used to inoculate 100 liters of medium M. The cells were grown aerobically at 37°C, overnight, until stationary. Five 100-liter fermentations were required to harvest 752 grams of wet cell pellet.

750 gram of *M. methylotrophus* cell pellet was suspended in 2.25 liters of Buffer A (20 mM Tris-HCl (pH 8.0), 50 mM NaCl, 1.0 mM DTT, 0.1 mM EDTA, 5% Glycerol) and passed through a Gaulin homogenizer at ~12,000 psig.



-23-

The lysate was centrifuged at  $\sim 13,000 \times G$  for 40 minutes and the supernatant collected.

5 The supernatant solution was applied to a 500 ml Heparin Hyper-D column (BioSeptra SA) which had been equilibrated in buffer A. A 1.0 L wash of buffer A was applied, then a 2 L gradient of NaCl from 0.05 M to 1 M in buffer A was applied and fractions were collected. Fractions were assayed for Mme I endonuclease activity  
10 by incubating with 1  $\mu$ g Lambda DNA (NEB) in 50  $\mu$ l NEBuffer 1, supplemented with 32  $\mu$ M S-adenosyl-L-methionine (SAM) for 15 minutes at 37° C. MmeI activity eluted at 0.3 M to 0.4 M NaCl.

15 The Heparin Hyper-D column fractions containing the Mme I activity were pooled, diluted to 50 mM NaCl with buffer A (without NaCl) and applied to a 105 ml Source15 Q column (Amersham Biotech) which had been equilibrated with buffer A. A 210 ml wash with buffer A was applied  
20 followed by a 1.0 L gradient of NaCl from 0.05 M to 0.7 M in buffer A. Fractions were collected and assayed from Mme I endonuclease activity. The Mme I activity was found in the unbound fraction.

25 The Source15 Q pool was loaded onto a 22 ml AF-Heparin-TSK column (TosoHaas) which had been equilibrated with buffer A. A wash of 44 ml buffer A was applied, followed by a linear gradient of NaCl from 0.05 M to 1.0 M in buffer A. Fractions were collected  
30 and assayed from Mme I endonuclease activity. The Mme I activity eluted between 0.26 M and 0.29 M NaCl. The fractions containing activity were pooled and dialyzed against buffer B (20 mM NaPO<sub>4</sub> (pH 7.0), 50 mM NaCl, 1.0 mM DTT, 0.1 mM EDTA, 5% Glycerol).

35

-24-

The dialyzed AF-Heparin-TSK pool was loaded onto a 6 ml Resource15 S column (Amersham Biotech) which had been equilibrated with buffer B. A wash of 12 ml buffer B was applied, followed by a linear gradient of NaCl from 0.05 M to 1.0 M in buffer B. Fractions were collected and assayed for Mme I endonuclease activity. Mme I activity eluted between 0.14 M and 0.17 M NaCl.

This pool was applied to a 2 liter Superdex 75 sizing column (Amersham Biotech) which had been equilibrated with buffer C (20 mM Tris-HCl, pH 8.0, 500 mM NaCl, 1.0 mM DTT, 0.1 mM EDTA, 5% Glycerol). Fractions were collected between 500 and 1500 ml elution with buffer C, then assayed by Mme endonuclease assay and polyacrylamide gel electrophoresis on 4-20% gradient gel, followed by protein staining with Coomassie Brilliant Blue dye. Fractions eluting between 775 and 825 ml corresponded to Mme I activity and a protein band of 105 kDa. These fractions were pooled and dialyzed against buffer D (20 mM NaPO<sub>4</sub>, (pH 7.0), 50 mM NaCl, 1 mM DTT, 5% Glycerol).

The dialyzed sizing pool was applied to a 16 ml Ceramic HTP column (BioRad) which had been equilibrated with buffer D. A 32 ml wash with buffer D was followed by a linear gradient from 0.02 M to 1.0 M NaPO<sub>4</sub> in buffer D. Fractions were collected and assayed by Mme endonuclease assay and polyacrylamide gel electrophoresis on a 4-20% gradient gel, followed by protein staining with Coomassie Brilliant Blue dye. Mme I eluted between 0.26 M and 0.3 M NaPO<sub>4</sub>. A portion of several fractions containing a single homogeneous protein band of 105 kDa were used for protein sequencing. The rest of the purified MmeI fractions were pooled (6 ml @ .36 mg/ml) and dialyzed against storage

-25-

buffer (10 mM Tris (pH 7.9), 50 mM KCl, 1mM DTT, .1 mM EDTA, 50% glycerol). The purified MmeI enzyme was stored at -20°C.

#### 5 Activity determination:

Samples from 1-4 µl were added to 50 µl substrate solution consisting of 1X NEBuffer 1, 32 µM S-adenosyl-L-methionine, and 1 µg DNA (lambda, PhiX174 or pUC19 DNAs). Reactions were incubated for 15 minutes at 37°, received 20 µl stop solution and were analyzed by electrophoresis on a 1% agarose gel.

#### 15 Optimized endonuclease activity

Following purification of MmeI from *M. methylotrophus*, experiments were performed to determine the optimal reaction conditions for DNA cleavage. Endonuclease activity was found to be significantly enhanced by the presence of potassium in the reaction buffer. Reactions were performed at 4°C to 37°C and from 5 to 60 minutes with no appreciable change in the amount of DNA cleavage. Enzyme concentrations at or near stoichiometric equivalence to DNA sites were required for maximal cleavage. Large excess of enzyme blocked cleavage. These findings were used to reassess the activity of MmeI and to define a workable endonuclease unit.

#### 30 Unit definition

One unit of MmeI is defined as the amount of MmeI required to completely cleave 1 µg of PhiX174 DNA in 15 minutes at 37°C in NEBuffer 4 (20 mM Tris-acetate, 10 mM magnesium acetate, 50 mM potassium acetate, 1 mM

-26-

dithiothreitol (pH7.9 at 25°C)) supplemented with 80 $\mu$ M S-adenosyl-L-methionine (SAM).

## EXAMPLE II

5

### CLONING THE *MmeI* ENDONUCLEASE

10

15

20

25

30

1. DNA purification: Total genomic DNA of *Methylophilus methylotrophus* was prepared. 5 grams of cell paste was suspended in 20 ml of 25% sucrose, 0.05 M Tris-HCl pH 8.0, to which was added 10 ml of 0.25 M EDTA, pH 8.0. Then 6 ml of lysozyme solution (10 mg/ml lysozyme in 0.25 M Tris-HCl, pH 8.0) was added and the cell suspension was incubated at 4°C for 16 hours. 25 ml of Lytic mix (1% Triton-X100, 0.05 M Tris, 62 mM EDTA, pH 8.0) and 5 ml of 10% SDS was then added and the solution incubated at 37°C for 5 minutes. The solution was extracted with one volume of equilibrated phenol:chloroform:isoamyl alcohol (50:48:2, v/v/v) and the aqueous phase was recovered and extracted with one volume of chloroform:isoamyl alcohol (24:1, v/v) two times. The aqueous solution was then dialysed against four changes of 2 L of 10 mM Tris, 1 mM EDTA, pH 8.0. The dialysed DNA solution was digested with RNase (100  $\mu$ g/ml) at 37°C for 1 hour. The DNA was precipitated by the addition of 1/10th volume 5 M NaCl and 0.55 volumes of 2-propanol and spooled on a glass rod. The DNA was briefly rinsed in 70% ethanol, briefly air dried and dissolved in 20 ml TE (10 mM Tris, 1 mM EDTA, pH 8.0) to a concentration of approximately 500  $\mu$ g/ml and stored at 4°C.

35

2. The *MmeI* endonuclease was purified to homogeneity as described in Example I above.

-27-

3. Amino acid sequences of the *MmeI* endonuclease were obtained for the amino terminus and for several internal cyanogen bromide digestion products of the *MmeI* polypeptide. The *MmeI* restriction endonuclease, prepared as described in Example I above, was subjected to electrophoresis and electroblotted according to the procedure of Matsudaira (Matsudaira. J. Biol. Chem. 262:10035-10038, 1987)), with modifications as previously described (Looney, et al. Gene 80:193-208 (1989)). The membrane was stained with Coomassie blue R-250 and the protein band of approximately 105 kD was excised and subjected to sequential degradation on an ABI Procise 494 Protein/Peptide Sequencer with gas-phase delivery (Waite-Rees, et al. J. Bacteriol. 173:5207-5219 (1991)). The amino acid sequence of the first 14 amino terminal residues obtained was the following:  
ALSWNEIRRKAIEF (SEQ ID NO:15).

An additional sample of the *MmeI* endonuclease, 20  $\mu$ g in 20  $\mu$ l, was treated with 2  $\mu$ g of cyanogen bromide (Sigma) dissolved in 200  $\mu$ l of 88% distilled formic acid for 24 hours in the dark at room temperature. This reaction mixture was evaporated to dryness and resuspended in 20  $\mu$ l of loading buffer (1.5M Tris-HCl, pH 8.5, 12% glycerol, 4% SDS, 0.05% Serva Blue G, 0.05% Phenol Red) at 100°C for 5 minutes. This sample was subjected to electrophoresis on a Tris-Tricine 10 to 20% polyacrylamide gradient gel (Invitrogen) for three hours and then transferred to a polyvinylidene difluoride (PVDF) membrane (Problott, Applied Biosystems Inc.) using 10 mM CAPS buffer (10mM 3-[cyclohexylamino]-1-propanesulfonic acid, 10% methanol, 0.05% SDS, 0.005% dithiotheritol, adjusted to pH 11.0 with NaOH) for 18 hours at 200 volts in a tank electroblotter (TE52, Hoeffer). The membrane was stained with Coomassie blue

-28-

R-250 and major bands of 25 kilodaltons (kD), 14 kD, 7.5 kD and 6 kD were observed, as well as smaller bands.

These stained protein bands were excised from the membrane and each subjected to sequential degradation.

5 The fragments other than the amino terminal fragment are derived from internal cleavage by cyanogen bromide at methionine residues from within the protein and thus should be preceded by a methionine. The first 29 residues of the 25 kD peptide corresponded to

10 (M)KISDEFGNYFARIPLKSTXXIXEXNALQ (SEQ ID NO:16). Residues 20, 21, 23 and 25, labeled X, were not identified. The first 40 amino acid residues obtained from the 14kD fragment were: (M)DAKKRRNLGAHYTSEANILKLI

15 KPLLLDELWVVFXXKVK (SEQ ID NO:17). Residue 36 was not determined. The first 25 residues of the 7.5 kD peptide corresponded to (M)KSRGKDLDKAYDQALDYFSGIAER (SEQ ID NO:18). The 6kD fragment was found to contain a mixture of three sequences.

20 4. Amplification of a portion of the MmeI endonuclease: The peptide sequence data from the amino terminus, 25 kD, 14kD and 7.5kD peptides was used to construct a series of degenerate PCR primers corresponding to the codons for the amino acid residues.

25 The order of the internal peptide fragments was unknown, so both forward (sense strand) and reverse (antisense strand) primers were made for these fragments. The primers were:

30 **25 kD fragment:** residues DEFGNYFA (SEQ ID NO:19)

Forward:

35 1) 5'-GARTTYGGNAAYTAYTTYGC-3' (SEQ ID NO:20)

Reverse:

2) 5'-AARTARTTNCCRAAYTCRTC-3' (SEQ ID NO:21)

-29-

**14 kD fragment:** residues MDAKKR (SEQ ID NO:22)

Forward A:

5 3) 5'-ATGGAYGCNAARAARCG-3' (SEQ ID NO:23)

Forward B:

10 4) 5'-ATGGAYGCNAARAARAG-3' (SEQ ID NO:24)

Reverse:

15 5) 5'-CGNCGYTTYTTNGCRTCCAT-3' (SEQ ID NO:25)

**7.5 kD fragment:** residues DKAYDQA (SEQ ID NO:26)

Forward:

20 6) 5'-GAYAARGCNTAYGAYCARGC-3' (SEQ ID NO:27)

Reverse:

25 7) 5'-GCYTGRTCRTANGCYTTRTC-3' (SEQ ID NO:28)

where

Y = T, C

R = A, G

H = A, T, C

30 S = G, C

N = A, C, G, T

Primers 1 and 2 are derived from the MmeI 25 kD CNBr peptide and were prepared to prime on the sense strand (1) or the antisense strand (2) of the gene. Primers 3 through 5 are derived from the 14 kD CNBr peptide and were prepared to prime on the sense strand (3 and 4) or the antisense strand (5) of the gene, with 3 and 4 differing in the codon usage for the arginine residue. Primers 6 and 7 are derived from the 7.5 kD CNBr peptide and were prepared to prime on the sense strand (6) or the antisense strand (7) of the gene.

45 PCR amplification reactions were performed using the primer combinations of 1 with 5, 1 with 7, 3 with 2,

-30-

3 with 7, 4 with 2, 4 with 7, 6 with 2 and 6 with 7. A portion of the MmeI gene was amplified in a PCR reaction by combining:

5        80  $\mu$ l 10X Thermopol buffer (NEB)  
      50  $\mu$ l 4mM dNTP solution (NEB)  
      4  $\mu$ l MmeI genomic DNA (500 $\mu$ g/ml stock)  
      16  $\mu$ l 100mM MgSO<sub>4</sub>  
      586  $\mu$ l dH<sub>2</sub>O  
10       16  $\mu$ l (32 units) Vent® exo- DNA polymerase (NEB).

      This master mix was divided into 8 aliquots of 90  $\mu$ l, to which was added 5  $\mu$ l forward primer (10  $\mu$ M stock) and 5  $\mu$ l reverse primer (10  $\mu$ M stock). The cycling  
15       parameters were 95°C for 3 minutes for one cycle, then 95°C for 30 seconds, 46°C for 30 seconds, 72°C for 2 minutes, for 25 cycles.

      The amplification reactions were electrophoresed on  
20       a 1% agarose gel and analyzed. Major DNA amplification products of 450 base pairs (bp) (primers 2 with 4), 650 bp (primers 5 with 6) and 1100 bp (primers 2 with 6) were obtained. These fragment sizes are consistent with the 7.5 kD CnBr fragment being located nearest the amino  
25       end of the protein and approximately 650 bp away from the 14kD CnBr fragment, with the 14 kD fragment between the 7.5 kD and the 25 kD fragment and adjacent to the 25 kD fragment. The amplified DNA fragments were gel  
30       purified and sequenced using the primers that were used for the amplification. A translation of the DNA sequence obtained matched the amino acid sequence derived from the purified MmeI endonuclease, indicating that a portion of the MmeI endonuclease gene DNA sequence had been successfully obtained.

35



-31-

5. Determining the DNA sequence for the entire MmeI gene and adjacent DNA: The inverse PCR technique was used to extend the DNA sequence from both sides of the 1060 bp of the MmeI gene obtained above. To accomplish this a series of primers matching the MmeI gene DNA sequence and oriented for inverse PCR were designed and synthesized. MmeI genomic DNA was cut with a number of restriction endonucleases and ligated at low concentration to generate circular DNA templates.

A. MmeI genomic DNA was digested with ten different restriction endonucleases and then circularly ligated to obtain DNA templates to amplify using the inverse PCR technique. The restriction enzymes used were:

BspHI (T/CATGA)  
EcoRI (G/AATTC)  
HindIII (A/AGCTT)  
HinfI (G/CGC)  
MspI (C/CGG)  
NlaIII (CATG/)  
PstI (CTGCA/G)  
SacI (GAGCT/C)  
SphI (GCATG/C)  
XbaI (T/CTAGA)

Restriction enzyme digests were performed by combining:

5 µl 10X NEBuffer recommended for the enzyme (varied with enzyme)  
2 µl *M. methylotrophus* genomic DNA (1 µg)  
43 µl dH<sub>2</sub>O  
1 µl (10 - 20 units) restriction enzyme.

The reactions were incubated for 1 hour at 37°C. The restriction endonuclease was inactivated by heating

-32-

the reaction to 65°C (80°C for PstI) for 20 minutes. The digested DNA was then ligated into circular fragments by adding 50 µl 10X T4 DNA ligase buffer, 400 µl dH<sub>2</sub>O and 3 µl concentrated T4 DNA ligase (6000 units, New England Biolabs, Inc.) and incubating at 16°C for 16 hours. The ligated DNA was then extracted with phenol and chloroform, precipitated with 2-propanol and resuspended in 100 µl TE buffer.

B. Amplification of DNA adjacent to the 1060 bp fragment of the MmeI endonuclease gene: Two pairs of PCR primers were designed, one near each end of the 1060 bp sequence obtained from direct PCR with degenerate primers. The primer sequences were:

primer IP 1:

5'-GTTGGATCCCGCACAGATTGCTCAGG-3' (SEQ ID NO:29)

primer IP 2:

5'-GTTGGATCCTACGTTAATCTGAATAAGATG-3' (SEQ ID NO:30)

primer IP 3:

5'-GTTGGATCCTGTTAATCTGAAACGCTGG-3' (SEQ ID NO:31)

primer IP 4:

5'-GTTGGATCCTTATACCAAATGTGAGGTC-3' (SEQ ID NO:32)

Inverse PCR reactions were performed on the 10 circularized templates produced above with the primer pairs of IP 1 with IP 2, IP 3 with IP 4, and IP 1 with IP 3. The amplification reactions were assembled by combining:

80 µl 10X Thermopol buffer (NEB)

50 µl 4mM dNTP solution (NEB)

40 µl IP primer (forward)

-33-

40 µl IP primer (reverse)  
16 µl 100mM MgSO<sub>4</sub>  
534 µl dH<sub>2</sub>O  
16 µl (32 units) Vent® exo- DNA polymerase (NEB).

5

10

The master mix was aliquoted into ten tubes of 76 µl, to which was added 4 µl of the appropriate digested, circularly ligated template. The cycling parameters were 95°C for 3 minutes for one cycle, then 95°C for 30 seconds, 56°C for 30 seconds, 72°C for 3 minutes, for 25 cycles. Amplification products were analyzed by agarose gel electrophoresis.

15

20

25

For primers IP 1 and IP 2 with the SphI template and the NlaIII template a product of approximately 825 bp was obtained. For primers IP 3 and IP 4 with the BspHI template a product of approximately 800 bp was obtained. For primers IP 1 and IP 3 with the EcoRI template a product of approximately 1500 bp was obtained. These amplified DNA fragments were gel purified, sequenced and assembled with that previously obtained. The assembled sequence did not contain the entire MmeI endonuclease open reading frame. The assembled sequence was used to direct synthesis of a second group of inverse PCR primer pairs. The sequences of these primers were:

primer IP 5:

30

5'-TTCAGAAATACGAGCGATGC-3' (SEQ ID NO:33)

primer IP 6:

35

5'-GTCAAGCCATAAACACCATC-3' (SEQ ID NO:34)

primer IP 7:

40

5'-GAGGGTCAGAAAGGAAGCTG-3' (SEQ ID NO:35)

-34-

primer IP 8:

5'-GTCCAACTAACCCTTTATGG-3' (SEQ ID NO:36)

5

10

15

20

25

Inverse PCR amplification reactions were performed as above. Using primers IP 5 and IP 6, products were obtained from the *Nla*III template (approximately 450 bp) and the *Msp*I template (approximately 725 bp), but not from the other circular ligation templates. Using primers IP 7 and IP 8, products were obtained from the *Eco*RI template (approximately 500 bp), the *Sph*I template (approximately 825 bp) and the *Bsp*HI template (approximately 750 bp). These DNA fragments were sequenced and the sequence was assembled with that previously obtained. The assembled sequence did not yet contain the entire *Mme*I endonuclease open reading frame, so another round of primer synthesis and inverse PCR was performed. Additional DNA templates were generated as above, but using the restriction enzymes *Apo*I (R/AATY), *Ase*I (AT/TAAT), *Bsa*HI (GR/CGYC), *Mfe*I (C/AATTG), *Ssp*I (AAT/ATT) and *Eco*RV (GAT/ATC) to digest *M. methylotrophus* genomic DNA. The sequences of this third round of primers were:

primer IP 9:

5'-TTCCTAGTGCTGAACCTTTG-3' (SEQ ID NO:37)

30

primer IP 10:

5'-GTTGCGTTACTTGAAATGAC-3' (SEQ ID NO:38)

35

primer IP 11:

5'-CCAAAATGGAACCTGTTTCG-3' (SEQ ID NO:39)

primer IP 12:

40

5'-GTGAGTGCGCCCTGAATTAG-3' (SEQ ID NO:40)

-35-

Inverse PCR amplification reactions were performed as above. Using primers IP 9 and IP 10, products were obtained from the NlaIII template (approximately 425 bp), the MfeI template (approximately 750 bp), the ApoI template (approximately 800 bp) and the MspI template (approximately 2100 bp). Using primers IP 11 and IP 12, products were obtained from the SphI template (approximately 875 bp), the BspHI template (approximately 925 bp) and the EcoRI template (approximately 950 bp). These DNA fragments were sequenced and the sequence was assembled with the sequences previously obtained. Further sequencing was performed on the IP 9, IP10 MspI 2100 bp product using three additional primers:

primer S1:

5'-GCTTCATTTTCATCCTCTGTGC-3' (SEQ ID NO:41)

primer S2:

5'-TAACCGCCAAAATTAATCGTG-3' (SEQ ID NO:42)

primer S3:

5'-CCACTATTCATTACAACACC-3' (SEQ ID NO:43)

The final assembled sequence (Figure 2) contained the entire MmeI restriction gene, as well as 1640 bp of sequence preceding the gene and 1610 bp of sequence following the gene.

6. Cloning the MmeI endonuclease gene in *E. coli*: The putative MmeI endonuclease open reading frame was identified from the DNA sequence assembly obtained

-36-

from sequencing the various inverse PCR amplified DNA fragments. The beginning of the open reading frame was identified on the basis of the match of the predicted amino acid sequence at the amino terminus of the open reading frame with the sequence determined from the MmeI endonuclease protein. The predicted end of the open reading frame would allow for the coding of an approximately 105 kD polypeptide, which matched the observed size of the native MmeI endonuclease. The amino acid sequence deduced from translation of this open reading frame contained conserved sequence motifs of N6mA DNA methyltransferases. However, no open reading frame containing sequence motifs conserved among DNA methyltransferases was observed adjacent to the MmeI endonuclease gene, as had been expected. It was decided to try to express the MmeI endonuclease in *E. coli* without having a second methyltransferase present to protect the *E. coli* host DNA from cleavage. Oligonucleotide primers were synthesized to specifically amplify the MmeI gene from *M. methylotrophus* genomic DNA for expression in the cloning vector pRRS (Skoglund, Gene 88:1-5 (1990)). The forward primer contained a PstI site for cloning, a stop codon in frame with the lacZ gene of the vector, a consensus *E. coli* ribosome binding site, the ATG start codon for translation (changed from the GTG used by *M. methylotrophus* to facilitate greater expression in *E. coli*) and 20 nucleotides that matched the *M. methylotrophus* DNA sequence:

5'-GTTCTGCAGTTAAGGATAACATATGGCTTTAAGCTGGAACGAG-3'  
(SEQ ID NO:44)

The reverse primer contained a BamHI site for cloning and 22 nucleotides that matched the *M.*

-37-

methylothrophus DNA sequence 3' to the end of the MmeI open reading frame:

5'-GTTGGATCCGTCGACATTAATTAATTTTTGCCCTTAG-3'

(SEQ ID NO:45)

The MmeI gene was amplified in a PCR reaction by combining:

10 50 µl 10X Thermopol buffer (NEB)  
30 µl 4mM dNTP solution  
12.5 µl forward primer (10µM stock)  
12.5 µl reverse primer (10µM stock)  
5 µl MmeI genomic DNA (500µg/ml stock)  
15 387 µl dH<sub>2</sub>O  
3 µl (6 units) Vent® DNA polymerase

20 The reaction was mixed and aliquoted into 5 tubes of 80 µl each. MgSO<sub>4</sub> was added (100mM stock) to bring the final concentration of Mg++ ions to 2mM, 3mM, 4mM, 5mM and 6mM respectively. The cycling parameters were 95°C for 30 seconds, 60°C for 30 seconds, 72°C for 3 minutes, for 24 cycles. The reactions were analyzed by gel electrophoresis and the 3mM through 6mM Mg++  
25 reactions were found to contain a DNA band of the desired size of 2.8kb. These reactions were pooled and the 2.8kb band was gel purified. The 2.8kb amplified MmeI gene fragment was digested with BamHI and PstI endonucleases (NEB) in the following reaction  
30 conditions:

15 µl 10X BamHI reaction buffer (NEB)  
1.5 µl BSA (NEB)  
50 µl MmeI gene 2.8 kb amplified DNA fragment  
35 80 µl dH<sub>2</sub>O

-38-

5  $\mu$ l BamHI endonuclease (100 units)

5  $\mu$ l PstI endonuclease (100 units)

5 The reaction was mixed and incubated for 1 hour at 37°C. The small fragments cleaved off the ends of the 2.8kb DNA fragment were removed, along with the endonucleases, by purification on a Qiagen QiaPrep spin column according to the manufacturer's instructions.

10 The cleaved MmeI gene DNA fragment was ligated to the pRRS vector as follows: 10  $\mu$ l of the digested, purified 2.8kb MmeI fragment was combined with 5  $\mu$ l pRRS vector previously cleaved with BamHI and PstI and purified, 5  $\mu$ l dH<sub>2</sub>O, 20  $\mu$ l 2X QuickLigase Buffer (NEB),  
15 the reaction was mixed, and 2  $\mu$ l of QuickLigase was added. The reaction was incubated at room temperature for 5 minutes. 5  $\mu$ l of the ligation reaction was transformed into 50  $\mu$ l chemical competent *E. coli* ER2683 cells and the cells were plated on L-broth plates  
20 containing 100  $\mu$ g/ml ampicillin and incubated at 37°C overnight. Approximately 200 transformants were obtained and 18 representatives were analyzed as follows: plasmid from each colony was isolated by miniprep procedures and digested with AlwNI and NdeI endonucleases to determine  
25 if they contained the correct size insert. 2 of the 18 transformants had the correct size insert of approximately 2800 bp. Both clones were tested to see if they produced MmeI endonuclease activity. The clones were grown overnight at 37°C in 500 mL L-broth  
30 containing 100  $\mu$ g/ml ampicillin. The cells were harvested by centrifugation, suspended in 10 mL sonication buffer (20mM Tris-HCl, 1mM DTT, 0.1mM EDTA, pH7.5) and broken by sonication. The crude lysate was cleared by centrifugation and the supernatant was  
35 recovered. The lysate was assayed for endonuclease



-39-

activity by serial dilution of the lysate in 1X reaction buffer NEBuffer 1 (New England Biolabs) containing 20 µg/ml lambda DNA substrate and supplemented with SAM at 100 µM final concentration. The reactions were incubated for 1 hour at 37°C. The reaction products were analyzed by agarose gel electrophoresis on a 1% agarose gel in 1X TBE buffer. One of the two clones had MmeI endonuclease activity. This active clone was designated strain NEB1457 and was used for subsequent production of MmeI. The plasmid construct expressing MmeI activity in this clone was designated pTBMmeI.1.

### EXAMPLE III

#### **THE MmeI ENDONUCLEASE PROVIDES IN VIVO PROTECTION AGAINST MMEI CLEAVAGE**

The plasmid pTBMmeI.1 was purified from NEB1457 using the Qiagen miniprep protocol. This plasmid has two MmeI sites in the vector backbone, and one site within the MmeI gene. The plasmid was digested with MmeI to test whether this DNA was resistant to MmeI endonuclease activity, which would indicate that the single MmeI gene was able to methylate DNA *in vivo* to protect the host DNA against its endonuclease activity. To test this the following were combined:

10 µl pTBMmeI.1 miniprep DNA  
15 µl 10X NEBuffer 4  
15 µl SAM (1mM stock solution)  
110 µl dH<sub>2</sub>O  
1 µl MmeI endonuclease (15 units)

The reaction was mixed and split in thirds. To one third was added 0.5 µl dH<sub>2</sub>O, to the second was added 0.5 µl pRRS vector and to the third was added 0.5 µl PhiX174

-40-

DNA as a positive control. The pTBMmeI.1 was not cleaved by the MmeI endonuclease activity, while the Phix174 and pRRS DNAs in the same reaction were cleaved, indicating that the three MmeI sites in the pTBMmeI.1 DNA are resistant to MmeI endonuclease activity (Figure 4).

#### EXAMPLE IV

##### **MmeI ENDONUCLEASE SENSITIVITY TO METHYLATION**

The prior literature reports that MmeI endonuclease methylates just one strand of its recognition sequence, and that this hemi-methylation does not block subsequent cleavage of the DNA by the endonuclease (Tucholski, Gene 223 (1998) 293-302). To test this a set of four oligonucleotides were synthesized so that a DNA substrate could be formed that was either unmethylated (oligo 1 + oligo 2), methylated in the top strand only (oligo 3 + oligo 2), methylated in the bottom strand only (oligo 1 + oligo 4), or methylated on both strands (oligo 3 + oligo 4). The oligos synthesized were:

Oligo 1:

5'-FAM-GTTTGAAGACTCCGACGCGATGGCCAGCGATCGGCGCCTCAGCTTT  
TG-3' (SEQ ID NO:46)

Oligo 2:

5'-FAM-CAAAAGCTGAGGCGCCGATCGCTGGCCATCGCGTCGGAGTCTTCA  
AAC-3' (SEQ ID NO:47)

Oligo 3:

5'-FAM-GTTTGAAGACTCCG(6mA)CGCGATGGCCAGCGATCGGCGCCTCAGCTT  
TTG-3' (SEQ ID NO:48)

-41-

Oligo 4:

5'-FAM-CAAAAGCTGAGGCGCCGATCGCTGGCCATCGCGTCGG (6mA)GTCTTCA  
AAC-3' (SEQ ID NO:49)

(Other nucleotides outside the MmeI recognition sequence were also methylated for other studies, but since MmeI does not have any sequence specificity for these nucleotides this does affect MmeI activity and these other methylations are omitted here for clarity.) Duplex DNA was formed by mixing 100 $\mu$ l top strand oligo (14 $\mu$ M stock) with 100 $\mu$ l bottom strand oligo (14 $\mu$ M stock), heating to 85°C and cooling slowly to 30°C over a time of 20 minutes. MmeI was then used to cleave the oligo pairs in a 30  $\mu$ l reaction of 1X NEBuffer4, 2.5  $\mu$ M oligo, 100  $\mu$ M SAM and 2.5 units MmeI. As a control, restriction endonuclease Hpy188I was also used to cleave the oligo DNA. The Hpy188I recognition sequence overlaps the first 5 nucleotides of the MmeI recognition sequence in this DNA, 5'-TCNGA-3' and is blocked by methylation at the adenine in either strand of the DNA. MmeI was found to cleave unmethylated DNA as expected. In contrast to previous teaching (Tucholski, Gene 223:293-302 (1998)) MmeI did not cleave the hemi-methylated DNA when the top strand only was methylated: 5'-TCCG(N6mA)C-3'. When the bottom strand only was methylated MmeI did cleave the DNA. When both strands were methylated MmeI did not cleave the DNA. (Figure 5) This finding is consistent with both the observed ability of the single MmeI enzyme to protect host DNA against cleavage in vivo and the observation that MmeI methylates only the top strand of its recognition sequence. We confirmed the report that MmeI enzyme methylates only the top strand of its recognition sequence by methylating the oligo

-42-

pairs above with tritium labeled H<sup>3</sup>-SAM, washing away the unincorporated SAM and counting the radioactivity in the DNA. Both the unmethylated oligo DNA and the top unmethylated, bottom methylated DNAs had greater than 10-fold more counts than background, while the bottom unmethylated, top methylated DNA and the DNA with both strands methylated had counts near background (Figure 6). These findings indicate that MmeI is a novel type of restriction modification system which does not require a separate methyltransferase enzyme to modify the host DNA to provide protection against the activity of the endonuclease, as is the case for the type IIG (also called type IV) enzymes such as Eco57I.

#### EXAMPLE V

##### **DNA SEQUENCING and ANALYSIS**

DNA Sequencing: DNA sequencing was performed on double-stranded templates on an ABI 373 or ABI 377 automated sequencer. Amplified DNA fragments and individual clones were sequenced with primers synthesized as above or from universal primers located in the vector.

Computer analyses: Computer analyses of the DNA sequences obtained were performed with the Genetics Computer Group programs (Deverex, et al., Nucleic Acids Res. 12:387-395 (1984)) and database similarity searches were performed via the internet at the National Center for Biotechnology Information site (<http://www.ncbi.nlm.nih.gov/BLAST/>) using the BLASTX and the BLASTP algorithms (Altschul, et al., J. Mol. Biol. 215:403-410 (1990) and Gish, et al., Nature Genet. 3:266-722 (1993)).

-43-

## WHAT IS CLAIMED IS:

5 1. Isolated DNA coding for the *MmeI* restriction enzyme, wherein the isolated DNA is obtainable from *Methylophilus methylotrophus*.

10 2. A recombinant DNA vector comprising a vector into which a DNA segment coding for the *MmeI* has been inserted.

15 3. Isolated DNA coding for the *MmeI* endonuclease and *MmeI*/methyltransferase, wherein the isolated DNA is obtainable from ATCC Accession No. PTA-4521.


4. Vectors that comprise the isolated DNA of claim 3.

20 5. A host cell transformed by the vector of claim 2 or 4.

25 6. A method of producing recombinant *MmeI* restriction endonuclease and *MmeI* methylase comprising culturing a host cell transformed with the vector of claims 2 or 4 under conditions suitable for expression of said endonuclease and methylase.

30 7. Isolated DNA coding for an *MmeI*-like restriction enzyme, wherein said isolated DNA hybridizes to at least one conserved motif of the nucleotide sequence coding for the *MmeI* restriction enzyme under predetermined conditions.

FIGURE 1: MmeI DNA CLEAVAGE



**FIGURE 2: MmeI DNA SEQUENCE**

1 GAATTC CAGA TAGGTAGTCC TTTGGTACTT CCATCCCAAC CAGTGT CACG  
51 TTCCGCGCCA AACCAATCGG TTAAAGTGTA AGAAAGTCTT GCACTGAAGT  
101 AGCTGTAGGA CAAACCGAAG TTAACCTCTG TGGTATCCCA GCGACCACCT  
151 TTAGGTGTTT GACGGAAGCC TGCTGCGTCA CCTGCCAAGT TATATTTCTT  
201 CCATGAACCA CCTGGGTACA GGTAGCTGAT CAAACCAGCA GTCCAACCCA  
251 AGCCTTCAAT AGCAGGAATA GTTCCGTTAT ACCCACCATA AATATCAATT  
301 TCGGCAGTTG CATCAGGGAA GGTATTTGGT GTCACGTTTG AACCCCATGC  
351 ACCGACATAA AAGCCGCTGT CATGAGTAAT ATCAATACCG CCTTGAACGG  
401 CAGGTTTGTG CCAGTTTGTG GAAATACCAC GAGCATAGTA ATCTGAAACA  
451 AATCCAACGT TTGCAGTAGC AGCCCAGGCT GATTTTTCTT CTTTAGCCTC  
501 TTCAGCTGCG TATGAAACTT GGGCAAAGA TAATGTGCTT AACACTGCTG  
551 TGAGCAATAT AGATTGACGC ATTAAGAGTC CTCCTCTCTGT GAAATCTTTG  
601 ATTAAGTTGT TGTAAACGAG AATGAAACAA CAACCACAAA GCAAAGCACG  
651 TGCCAAACTA TAAATAACAT TATAATCAAT TATTTAAAAT ATATTTATAA  
701 TCTAAAATAT TAAATTAATT ATTTAATAAA CTGTTTTTTA TTGATTTAAC  
751 TCTAAAACAT ATGGGTGCAA CCACCCTTTT TACTCACTGA TAATGCTAAN  
801 ATAGCCAACA AAGGAGCCTT CACCATGCTG ATTTCAAATG AAAAAATTCA  
851 GGAATTATCT TTAAAAATCA AACAACTAAT CGAATCAAGC CCCATTTTCAG  
901 AGCTAAATAA CAACTTGCAT GCACTAATTC AGGGCGCACT CACCAAATG  
951 GAACTTGTTT CGCGTGAAGA ATTCGATATC CAATCTGCAT TATTAGCGCG  
1001 CACGCAAGAG CAATTAAAAC GTCTTGAAGA AAAAATCAGC CAGCTTGAAG  
1051 AAGGGCAGGC ATCCAGAAAG TAAAAATTAA TTTACAATTG TTAGCATTCC  
1101 ATTATTGAGG AGTGCGCTAT GAGTCTGGCG GTGTTATACA GTCGCGCGTT

1151 AAGCGGCATG GAGGCGCCAG AAGTGGTGGT AGAAGTCCAC TTGGCGAATG  
1201 GACTACCCAG CTTTACCATT GTTGAAACAT ATTGAAACTT TAAGCCTTAG  
1251 CATTTTTTCA AATATACAAA TGCCCCAAGC TGGTGCATTA AGAAGAATGT  
1301 AACAACTCCC TGCAGACTAG GAATAACTTC ATGATTTAAC GAACATCCCT  
1351 GAGTTTCAAA GTCGAATCTT CTCGTGTTGC AAATTTCTAC AGCTTCCTTT  
1401 CTGACCCTCT TGCACCAAAT TGCACATATGG CGCTAATAAA TCTTCTGCTA  
1451 TCCAATAATG TCCAACATAAC CCTTTATGGA CTCTPAAAAA AGATTTAATA  
1501 AATGATTAAAG ATGAATTCAA GGAATTTGAT GCCTGGAAAAT ATGGCAAAAG  
1551 CAAAAAGGCA GCCCAGTGCT GACTTTTTTG TTTTAACATT GGCCCATATA  
1601 TCCAATTTCA AATAATTTAA AAATTATCGG GAGCTAATCT GTGGCTTTAA  
1651 GCTGGAACGA GATAAGAAGA AAAGCTATTG AGTTTCTTAA AAGATGGGAA  
1701 GACGCCTCAG ATGAAAACAG TCAAGCCAAA CCCTTTTTAA TAGATTTTTT  
1751 CGAAGTTTTT GGAATAACTA ATAAGAGAGT TGCAACATTT GAGCATGCTG  
1801 TCAAAAAGTT CGCCAAGGCC CATAAGGAAC AATCTCGAGG ATTCTAGAT  
1851 TTGTTTTGGC CTGGCATTCT TCTTATTGAA ATGAAAAGCA GAGGTAAAGA  
1901 CCTCGACAAA GCGTATGACC AGGCACTTGA TTACTTTTCT GGCATTGCAG  
1951 AAAGAGACTT ACCCAGATAC GTTTTAGTTT GCGACTTCCA GCGTTTCAGA  
2001 TTAACAGACC TAATAACAAA AGAGTCAGTT GAATTTCTTT TAAAGGACTT  
2051 ATACCAAAAT GTGAGGTCTT TTGGTTTTAT AGCTGGTTAT CAACTCAAG  
2101 TAATCAAGCC ACAAGACCCT ATTAATATTA AGGCGGCTGA ACGGATGGGT  
2151 AAGCTTCATG ACACCCTGAA GTTGGTTGGA TATGAGGGAC ACGCTTTAGA  
2201 ACTTTATCTA GTGCGTTTAC TTTTTTGCTT ATTCTGAGAA GACACAATA  
2251 TTTTGTAGAA AAGTTTATTC CAAGAATATA TCGAGACAAA GACGCTAGAG  
2301 GACGGCAGTG ACCTTGCACA TCATATCAAT AACTTTTTT ATGTTCTCAA  
2351 TACCCAGAA CAAAAAGAT TAAAGAATCT AGACGAACAC CTTGCTGCAT



2401 TTCCATATAT CAATGGAAAA CTTTTCGAGG AGCCACTTCC GCCAGCTCAG  
2451 TTTGATAAAG CAATGAGAGA GGCATTGCTT GACTTGTGCT CATTAGATTG  
2501 GAGCAGGATT TCACCAGCAA TATTTGGAAG TTTATTCCAA AGCATTATGG  
2551 ATGCTAAAAA GAGAAGAAAT CTTGGGGCAC ACTACACCAG CGAAGCAAAT  
2601 ATTCTCAAGT TAATCAAGCC ATTGTTTCTT GACGAGCTCT GGGTAGAGTT  
2651 CGAGAAAGTT AAAAATAATA AAAATAAATT ACTAGCGTTC CACAAAAAAC  
2701 TAAGAGGACT TACATTTTTC GACCCTGCAT GCGGTTGCGG AAATTTTCTT  
2751 GTAATCACAT ACCGAGAACT AAGACTTTTA GAAATTGAAG TGTTAAGAGG  
2801 ATTGCATAGA GGTGGTCAAC AAGTTTGGGA TATTGAGCAT CTTATTCAGA  
2851 TTAACGTAGA CCAGTTTTTT GGTATCGAAA TAGAGGAGTT TCCCGCACAG  
2901 ATTGCTCAGG TTGCTCTCTG GCTTACAGAC CACCAAATGA ATATGAAAAT  
2951 TTCAGATGAG TTTGGAAACT ACTTTGCCCC TATCCCCTA AAATCTACTC  
3001 CTCACATTTT GAATGCTAAT GCTTTACAGA TTGATTGGAA CGATGTTTTA  
3051 GAGGCTAAAA AATGTTGCTT CATATTAGGA AATCCTCCAT TTGTTGSETA  
3101 AAGTAAACAA ACACCGGGAC AAAAAGCGGA TTTACTATCT GTTTTTGGAA  
3151 ATCTTAAATC CGCTTCAGAC TTAGACCTAG TTGCTGCTTG GTATCCCAAA  
3201 GCAGCACATT ACATTCAAAC AAATGCAAAC ATACGCTGTG CATTTGTCTC  
3251 AACGAATAGT ATTACTCAAG GTGAGCAAGT ATCGTTGCTT TGGCCGCTTC  
3301 TGCTCTCATT AGGCATAAAA ATAACTTTG CTCACAGAAC TTTCAGCTGG  
3351 ACAAATGAGG CGTCAGGAGT AGCGGCGGTT CACTGCGTAA TTATCGGATT  
3401 TGGGTTGAAG GATTCAGATG AAAAAATAAT CTATGAGTAT GAAAGTATTA  
3451 ATGGAGAACC ATTAGCTATT AAGGCAAAAA ATATTAATCC ATATTTGAGA  
3501 GACGGGGTGG ATGTGATTGC CTGCAAGCGT CAGCAGCCAA TCTCAAAATT  
3551 ACCAAGCATG CGTTATGGCA ACAAACCAAC AGATGATGGA AATTTCTTAT  
3601 TTAGTGACGA AGAAAAAAC CAATTTATTA CAAATGAGCC ATCTTCCGAA


3651 AAATACTTCA GACGGTTTGT GGGCGGGGAT GAGTTCATAA ACAATACAAG  
3701 TCGATGGTGT TTATGGCTTG ACGGTGCTGA CATTTTCAGAA ATACGAGCGA  
3751 TGCCTTTGGT CTTGGCTAGG ATAAAAAAG TCCAAGAATT CAGATTAAAA  
3801 AGCTCGGCCA AACCAACTCG ACAAAGTGCT TCGACACCAA TGAAGTTCTT  
3851 TTATATATCT CAGCCGGATA CGGACTATCT GTTGATACCT GAAACATCAT  
3901 CTGAAAACAG ACAATTTATT CCAATTGGTT TTGTTGATAG AAATGTCATT  
3951 TCAAGTAACG CAACGTATCA TATTCCTAGT GCTGAACCTT TGATATTTGG  
4001 CCTGCTTTCA TCGACCATGC ACAACTGCTG GATGAGAAAT GTAGGAGGAA  
4051 GGTTAGAAAG TCGTTATAGA TATTCTGCCA GCCTGGTTTA CAACACGTTT  
4101 CCATGGATTC AACCCAACGA AAAACAATCG AAAGCGATAG AAGAAGCTGC  
4151 ATTTGCGATT TTAAGGCTA GAAGCAATTA TCCAAACGAA AGTTTAGCTG  
4201 GTTTATACGA CCCAAAAACA ATGCCCTAGTG AGCTTCTTAA AGCACATCAA  
4251 AAACCTGATA AGGCTGTGGA TTCTGTCTAT GGATTTAAAG GACCAAACAC  
4301 ACAAATTGCT CGAATAGCTT TTTTGTTTGA AACATACCAA AAGATGACTT  
4351 CACTCTTACC ACCAGAAAAA GAAATTAAGA AATCTAAGGG CAAAAATTAA  
4401 TTAATGTATT TAACATTAAA CCACCCTGAT TTATTTTCGAA TAGTTCAAAT  
4451 GCTTCCATGT GGAATAATCG CCTTCAATCA TATTAAAAAA CCGACGCTAG  
4501 TAATAAAAAC TTCCAAAGAG GCCATATTAA CCGCCAAAAT TAATCGTGAA  
4551 TTTAAATAT ATCTTTATCA AACCACATCG GCTTGTGTTC TAGTAAGTGC  
4601 ATTTTTTGAC GATTCTGATA GTCCACTATT CATTACAACA CCAATTGTTC  
4651 GAGATGACCA AACTCCTTA GACTTGTTAA GATTTTAAAT CAACAATGAT  
4701 TTTACGATTT GCTTCTTTGA TGAAGTGAAC CGAGAATTTT TTTCCGTAA  
4751 CGCAACTGGT AATTTAGTCT CTATCTTTGA GAGCATTAC TTGATGCCAC  
4801 TGCCGAGCCC AGAGGAAGCC CACAATGCAT TGAATGAAGC GGAATTTTGG  
4851 TTCAGTTTAC GCTCAGCTGC TGATGATGAA TCATCTATCC AGGTTTCTTT

4901 ATTGGATAAT CTATTTCTTG ACGATTTTGT AATTTATGAC CTATCCTCAA  
4951 ACAAAAACGA TATGACATCA TTGGTTAGAG AAATAAACC AGGATACTAT  
5001 CAGGAAGCAG ATATTGCAAA GTTACTAACA AGAGCTTTTA GTTTGGAAAG  
5051 CATTTATCAG AATCCAGTGA AAACAAGCGA TTCAAAAGAG TTGGCAGACG  
5101 TTGTGGTATT CGGCCAAAAG GAAATTTTAA TAATTCAAGC TAAAGATAGT  
5151 GAAAACAATC AGAAACAAGT TTTAGAGGTT TCGTTAGACA AGAAATGCGC  
5201 AAAGTCTTCA AAGAACTTT CTGAAGCTTT GGCACAACCTC ACCGACACTA  
5251 TCTTAACAAT ATCCAATACA CCAATAGTTG ATGTTCCGGT TGGTAAGAAA  
5301 AAATGCACTC TGAACCTTGA GGGAAAGCAG CTTATTGGTA TCGTCGTTGT  
5351 TAAAGAGCTT TTTAATGATA TTTACGATAA ATACAGTCAA AAAGTTTTTG  
5401 AGCATGTAGA GTTGTCTAAA GCACCCATTG TCTTCTTTGA CTATCCAGAA  
5451 TTTGCAAGAA TGACATTTCA TTGTAATTCT GAGGAATTAT TACTTTATGC  
5501 TTTGCATAGG ATATTTAGTT CTGCAATAGA AAATGGAATG TATAAACGAT  
5551 TGAGATTTAC TCAACCTATC ATAACTGATG GTCATGACAG CTACTTCAGG  
5601 ATACAAAACA GGCCCCATTC TGATGAGGCC TATTTAATTT GCACAGAGGA  
5651 TGAAATGAAG CTCTCAAATA AGTTTAAAGA CTAAATTTAT ATTTTCCTCA  
5701 GTATCTTAAA AACAATATTC ATTAAATTGG AAAGCCCGCA ATGATTGTTG  
5751 CAGTATCAAT GCGGGCATCA GTATCCAGCT CTTGCAATAC ACGGAAGTAT  
5801 CAAGAAGCGA ATCAGGATTC TAACCATAACC TTTTAAATTG CAACAATCTA  
5851 ATTTCCATAA CATGTGTAGC TACATCGAAA AAAAGACCTC GAAGAGGTTG  
5901 CAAGAGCGTC CAGCTCGCGG CATCAAAAGA CCCTAGTCTT TTGACAAGGG  
5951 GGAGCCAAAA AACTGAGGTG GAGGAGCTTG CCGACGAAGC CAGGAAGCCC  
6001 CAGCGTCCGG

**FIGURE 3: MmeI AMINO ACID SEQUENCE**

1 MALSWNEIRR KAIEFSKRWE DASDENSQAK PFLIDFFEVF GITNKR VATF  
51 EHAVKKFAKA HKEQSRGFVD LFWPGILLIE MKSRGKDLDK AYDQALDYFS  
101 GIAERDLPRY VLVCDFQRFR LTDLITKESV EFLLDLYQN VRSFGFIAGY  
151 QTQVIKPQDP INIKAAERMG KLHDTLKLVG YEGHALELYL VRLLFCLFAE  
201 DTTIFEKSLF QEYIETKTLE DGSDLAHHIN TLFYVLNTPE QKRLKNLDEH  
251 LAAPFYINGK LFEEPLPPAQ FDKAMREALL DLCSLDWSRI SPAIFGSLFQ  
301 SIMDAKKRRN LGAHYTSEAN ILKLIKPLFL DELWVEFEKV KNNKNKLLAF  
351 HKKLRLGLTFF DPACGCGNFL VITYRELRL L EIEVLRGLHR GGQQVLDIEH  
401 LIQINVDQFF GIEIEEFPAQ IAQVALWLTD HQNMNKISDE FGNYFARIPL  
451 KSTPHILNAN ALQIDWNDVL EAKKCCFILG NPPFVGKSKQ TPGQKADLLS  
501 VFGNLKSASD LDLVAAWYPK AAHYIQTNAN IRCAFVSTNS ITQGEQVSL L  
551 WPLLLSLGIK INFARTFSW TNEASGVA AV HCVIIGFGLK DSDEKIIY EY  
601 ESINGEPLAI KAKNINPYLR DGVDVIACKR QQPISKLP SM RYGNKPTDDG  
651 NFLFTDEEKN QFITNEPSSE KYFRRFVGGD EFINNTSRWC LWLDGADISE  
701 IRAMPLVLAR IKKVQEFRLK SSAKPTRQSA STPMKFFYIS QPDTDYLLIP  
751 ETSSENRQFI PIGFVDRNVI SSNATYHIPS AEPLIFGLLS STMHNCWMRN  
801 VGGRLESRYR YSASLVYNTF PWIQPNEKQS KAIEEAAFAI LKARSNYPNE  
851 SLAGLYDPKT MPSELLKAHQ KLDKAVDSVY GFKGPNT EIA RIAFLFETYQ  
901 KMTSLLPPEK EIKKSKGKN\*


FIGURE 4: pTBMmeI.1 IS RESISTANT TO MmeI CLEAVAGE



9/17

FIGURE 5: MmeI CLEAVAGE OF HEMI-METHYLATED SUBSTRATES

1 2 3 4 5 6 7 8 9 10 11 12 13 14



10/17

**FIGURE 6: METHYLATION INCORPORATION BY MmeI ENDONUCLEASE**

Top Strand:	/	Bottom strand:	<sup>3</sup> H-COUNTS
5'-TCCG <u>A</u> C-3'	/	5'-GTCGGA-3'	
unmethylated:	/	unmethylated	19,972
unmethylated:	/	methylated:	14,447
methylated:	/	unmethylated:	1,266
methylated:	/	methylated:	917

'A' indicates position of N6-methyl adenine in the DNA substrate

## MmeI Figure 7: Multiple Sequence Alignment of MmeI and Homologous Polypeptides

PileUp of: @mme.list2

Symbol comparison table: GenRunData:blosum62.cmp CompCheck: 1102

GapWeight: 6

GapLengthWeight: 1

Name: mmelfeLORF3P = gi|28373198|ref|NP\_783835.1| (SEQ ID NO:3)  
 Name: mmeLre121P = gi|23451826|gb|AAN32874.1|AF461726\_1 (SEQ ID NO:4)  
 Name: mme = MmeI amino acid sequence (SEQ ID NO:5)  
 Name: mmeNMA1791 = gi|15794682|ref|NP\_284504.1| (SEQ ID NO:6)  
 Name: mmeBSU0677 = gi|16077744|ref|NP\_388558.1| (SEQ ID NO:7)  
 Name: mmeGcry = gi|9945797|gb|AAG03371.1| (SEQ ID NO:8)  
 Name: mmePflQ8 = gi|23451826|gb|AAN32874.1|AF461726\_1 (SEQ ID NO:9)  
 Name: saro3834 = gi|23110638|gb|ZP\_00096791.1| (SEQ ID NO:10)  
 Name: mmeMSI135 = gi|20803963|emb|CAD31540.1| (SEQ ID NO:11)  
 Name: mmeCC0826 = gi|16125079|ref|NP\_419643.1| (SEQ ID NO:12)  
 Name: mmeDR0119.1 = gi|15807788|ref|NP\_285443.1| (SEQ ID NO:13)  
 Name: mmeDR2267 = gi|15807258|ref|NP\_295988.1| (SEQ ID NO:14)

	1				50
mmelfeLORF3P	-----	-----MPT	RQQAAREFVK	TWS.SDKKGR	EDADRQTFWN
mmeLre121P	-----	-----	-----	-----	-----
mme	-----	---MALSWNE	IRRKAIEFSK	RWE.DASD..	ENSQAKPFLI
mmeNMA1791	-----	---MKTLLO	LQTAAQNFAA	YYK.DQTD..	ERREKDTF*N
mmeBSU0677	-----	---MALID	LEDKIAEIVN	R.E.DHSD..	.....FLY
mmeGcry	MVMAPTTFVD	RATIRHNLTE	FKLRWLDRIK	QWEAENRPAT	ESSHDQQFWG
mmePflQ8	-----	-----	-----	-----	-----
saro3834	-----	-----	-----	-----	-----
mmeMSI135	-----	---MSLGAAGL	TPITPAAFIK	KWRKSELG..	ERQAAQEHFL
mmeCC0826	-----	-----	---MTPAQFVK	KWSDSQLR..	ERQASQEHFL
mmeDR0119.1	-----	-----	---MHPQEFAD	TWSRRALKAT	ERDSYVQHWL
mmeDR2267	-----	---MPQTE	TAQRMEDFVA	YW..RTLKGD	EKGESQV.FL
	51				100
mmelfeLORF3P	DLQVRVYDID	N.YYDYITYE	KDVQVKADGK	VTTRRIDGYI	P.STKIMVEM
mmeLre121P	-----	-----	-----	-----	-----
mme	DFFE.VFGIT	N.KR...VAT	FEHAVKKFAK	AHKEQSRGFV	DL...FWPGI
mmeNMA1791	EFFA.IFGID	R.KN...VAH	FEYPVKD..P	ADNTQ...FV	DI...FWEGI
mmeBSU0677	ELLG.VYDVP	R.AT...ITR	....LKK.GN	QNLTKRVGEV	HLKNKVV...
mmeGcry	DLDC.FGV.	N.ARDLYLY.	....QSAK	RASTGRTGKI	DM...FMPGK
mmePflQ8	-----	-----	-----	-----	-----
saro3834	-----	-----	-----	-----	-----M
mmeMSI135	D.ICSLVGHP	SP.SDEDPTG	AFFAFEKGAN	KLK.GGKGFA	D.VWK..KGH
mmeCC0826	D.LCRMLEVP	TP.AEDDPLG	ERYCFERGAA	KTG.GGDGWA	D.VWR..KGC
mmeDR0119.1	D.LCQLLHHE	APGADPD...	..YKFERRVT	KVGTKDKGFA	D.VFK..KAH
mmeDR2267	DRLFQAFGH.	...AGYKEAG	AE..LEYRVA	KQG.GGKKFA	DLLWR...PRV
	101				150
mmelfeLORF3P	KGKNIKDLK	PITQSGGD..	.....ELT	PFEQAKRYAN	FLPN...SEQ
mmeLre121P	-----	-----	-----	-----	-----
mme	LLIEMKSRGK	DL.....D..	.....KAY	D..QALDYFS	GIAERD...L
mmeNMA1791	FLAEHKSANK	NL.....T..	.....KAK	E..QAERYLQ	EIGRTKPSAL
mmeBSU0677	.FKKAK.KGK	LF.....D..	.....ALI	DIEQQVEYL.	.....SAK





saro3834	IFVGEGLFSR	TVETMSARDA	SDTHMVIAEI	FRAMDTRLAD	RAAAGIKSWA
mmeMSI135	LLPD.GLFTK	LLK.RSARAP	ERAMSYLDKL	FEAME.....	RGGEF...DL
mmeCC0826	LLPE.GLFTR	LTRSMQMRPP	AEAAPQFDAL	FAMMR.....	AGGMF...GA
mmeDR0119.1	MLER.GIVTR	LLE.RARAPP	GEDQLYFQDL	FGAMK.....	GGGEF...WG
mmeDR2267	LIPR.GFFTE	LADD.ARAGR	GSSFDLFGGL	FRQMTSERA	RGGRF.....
	351				400
mmelfeLORF3P	NQFAYVNGGM	FSDENVIIPQ	FTDELKRLIV	EDAGRGFDWS	GISPTIFGAV
mmeLre121P	-----	-----	-----	-----	-----
mme	AAFPYINGKL	FEEPLPPA.Q	FDKAMREALL	.DLCS.LDWS	RISPAIFGSL
mmeNMA1791	KGFEYVNGGL	FKERLRTF.D	FTAKQHRALI	.DCGN.FDWR	NISPEIFGTL
mmeBSU0677	KEFPYVNGQL	FTEPHTEL.E	FSAKSRKLII	.ECGELLNWA	KINPDIFGSM
mmegcry	AKFPYVNGAL	FAEPLAS.EY	FDYQMREALL	AAC..DFDWS	TIDVSVFGSL
mmePflQ8	-----	-----	-----	-----	-----
saro3834	DVFPYVNGQL	FSGS.TECPR	FSKIARSYLL	H..IGSLDWQ	KINPDIFGSM
mmeMSI135	TDITWFNGGL	FDGR..RALR	LDDGDIGLL.	.VAADSLDWG	LIDPTIFGTL
mmeCC0826	DIVHWFNGGL	FDEK..PALP	LERADIKLIH	DTAAEH.DWS	DLDPVFGNM
mmeDR0119.1	TDIRHFNGGL	FDSE..DALA	LTSEDAAAL.	.IIAAKLDWS	EVEPSIFGTL
mmeDR2267	APIPYFNGGL	FRAV..DPIE	LNRDELYLLH	KAALEN.NWA	RIQPQIFGVL
	401				450
mmelfeLORF3P	FESTLN.PET	RRSGGMHYTS	IENIHKVIDP	LFLNDLHDEF	D.....
mmeLre121P	-----	-----	-----	-----	-----
mme	FQSIMD.AKK	RRNLGAHYTS	EANILKLIK	LFLDELWVEF	E.....
mmeNMA1791	FQSVMD.AQE	RREAGAHYTE	AANIDKVIING	LFLLENLRAEF	E.....
mmeBSU0677	IQAVAS.EES	RSYLGMYHYS	VPNIMKVIK	LFLDKLNQSF	.....
mmegcry	FQLVKS.KEA	RRSDGEHYTS	KANIMKTIGP	LFLDELRAEA	D.....
mmePflQ8	-----	-----	-----	-----	-----
saro3834	IQAVAD.DEE	RGALGMHYTS	VPNILKVLNP	LFLDDLRAKL	E.....
mmeMSI135	FERFLD.PEK	RAQIGAHYTD	PEKIMRLVDP	VILRPLRQEW	EQARREIVEL
mmeCC0826	FEEALKATRE	RAALGAHYTD	REKILKIIDP	VITWPLMAQW	ETALAEIRAA
mmeDR0119.1	FENSLDV.DT	RSRRGAHYTS	VNDIERIVDR	VVMEPLWAEW	D.....
mmeDR2267	FQSSMDKKEQ	HAK.GAHYTS	EADIMRVVLP	TIVTPFQRQI	EAATTQ....
	451				500
mmelfeLORF3P	.....	.....	.....	.....	....KIQNMG
mmeLre121P	-----	-----	-----	-----	-----
mme	.....	.....	.....	.....	....KVK...
mmeNMA1791	.....	.....	.....	.....	....AVK..A
mmeBSU0677	.....	.....	.....	.....	.....L
mmegcry	.....	.....	.....	.....	....KL..VS
mmePflQ8	-----	-----	-----	-----	-----
saro3834	.....	.....	.....	.....	....E.....
mmeMSI135	LNGN.....	.....	..RKPPMRR.	..QOSRR...	....MKREEAA
mmeCC0826	LDARAAAEAE	RKAVLEAAAE	AMRADPVKAK	AGEAARRKTL	TAIAKRSDA
mmeDR0119.1	.....	.....	.....	....ALRLSL	PELKK.....
mmeDR2267	.....	.....	.....	.....	.....
	501				550
mmelfeLORF3P	NRRQRVTRAK	AFRDKLGKLK	FFDPACGSGN	FLTETYLSLR	KMENECLRII
mmeLre121P	-----	-----	-----	-----	-----
mme	...NNKNKLL	AFHKKLRLGT	FFDPACGCGN	FLVITYRELR	LLEIEVL.RG
mmeNMA1791	LKRDKAKKLA	AFYQKIQNLQ	FLDPACGCGN	FLIVAYDRIR	ALEDDIIAEA
mmeBSU0677	DAYDDYTKLE	NLLTRIGKIK	FFDPACGSGN	FLIITYKELR	RMEINI IKRL
mmegcry	SPSTSVAAL	RFRDSLSELV	PADMACGSGN	FLLLAYRELR	RIETDIIVAI
mmePflQ8	-----	-----	-----	-----	-----
saro3834	.AGDNSRKLL	NLRNRMKIR	VFDPAACGSGN	FLVIAYKQMR	ELEAEI....
mmeMSI135	.AEVR.SR..	.FTELRKLR	ILDPACGSGN	FLYLALQGVK	DIEHRANLDC
mmeCC0826	LGQAK..DRLE	AFLSRLAFA	VLDPAACGSGN	FLYVFLHALK	DIERRALVDA

```

751
mmelfeLORF3P VLFNDFHIFI NYGYRSFEWN NEAANKAKVD VVIVGFSTK. .EDKNPTIYD 800
mmelre121P ~~~~~ -VNKDKILYN
mmelre121P LLLS.LGIKI NFAHRTFSWT NEASGVAAVH CVIIGFGLKD ..SDEKIIYE
mmelre121P SLLN.QGIEI HFAHRTFQWT SQAAGKAAVH CIIVGFRQKP PMPSEKTLVD
mmelre121P ELFK.FGIQI NFAYKSFKWA NNAKNNAAVI VVIVGFG... PLDTKVNKYL
mmelre121P PIFKA.GWRI RFAHRTFAWD SEAPGKAAVH CVIVGFDKES ..QPRPRLWD
mmelre121P ~~~~~
mmelre121P RILKSAN..V KFAYRPFWRWS NSAANNAGVY CTIIGLTGSE VSNKK.....
mmelre121P RIIA.ES.RL FEAWSDEPWV VDG...AAVR VSLICFGHG. .EDPLCL...
mmelre121P PIAD.AG.AL MEAWADEPWA LEG...AAVR VSMFGFGDG. .FAERRL...
mmelre121P RIIA.TG.DL FMAWPDEPWQ QNG...AAVR VSLFGFDNG. .TETLRT...
mmelre121P YVVQHGG.TI TDAVGTQVWS GD....AAVH VSIWNWVKGP AEGPKHLAWQ

801
mmelfeLORF3P EQKIIS.... ..A.KHINQY MYSDNIFID TTRKY.IEA. MPKMKGTGNRP 850
mmelre121P SSN*ISH... ..C.KNINGY LFDGNMIFV. TNRPAPLSN. VPRMHNGCKL
mmelre121P YESINGEPLA IKA.KNINPY LRDGVDVIA. CKRQQPISK. LPSMRYGNKP
mmelre121P YPDIKGEPEK HAV.ANINPY LIDAPDLII. AKRSRPIHC. EPDMVNGSKP
mmelre121P FVD...ETKK L.V.SNISPY LTDGENILV. SSRTPKISD. LPKLHFGNMP
mmelre121P YPDVKGEVVS VEVGQSINAY LVDGPNVLVD KSR.HPISSE ISPATFGNMA
mmelre121P ~~~~~
mmelre121P ...LFGEGSV VEC.SSIAPY LVPGEPI.IC APRQSSIS.G FARMVMGNSP
mmelre121P ..DGRT.... ..VAQINAD LTAGVTDLTK ARRL.S..EN QNVAFMGDTK
mmelre121P ..EGRK.... ..AEHLHSD FRGASTDVTQ ALRL.K..EN ASIAFMGDTK
mmelre121P LNDGH..... ..VGVINAD LNAG.TDVKQ AQKL.P..EN AGVSFIGTQK
mmelre121P VGDHRTSPWQ STELPVINS A LSAG.TDVTQ AQKL.RVNMM SGACYQGQTH

851
mmelfeLORF3P ADGGALILSP KEAKEL.VNE EPQSKQ...F IKKLTGSKEF IT.GKYRYCL 900
mmelre121P LDGGFYTLTS QERKEA.ISK DPYADK...F IRPYLGAKNF IH.GTARYCI
mmelre121P TDDGNFLFTD EEKNQF.ITN EPSSEK...Y FRRFVGGDEF IN.NTSRWCL
mmelre121P TEGGNLILST AEKDAL.IAA EPLAEQ...Y IRPFIGADEF LN.GKTRWCL
mmelre121P NDGGGLLFTI TEYTD.AINK YPELVP...Y FKKFIGSVEF IN.GGLRYCL
mmelre121P RDGGNLLVEV DEYDEV.MS. DPVAAK...Y VRPFRGSREL MN.GLDRWCL
mmelre121P VEGGNLLFAE EEKQRL.VEG NVDVVK...F LKRUYGASEY IR.GEVRFCL
mmelre121P VDGKRLIFEQ DEKESV.VAA DPRSER...F FKRYGGTQEL VN.GVDRWCL
mmelre121P G.GAFDVPGS LARAWLSMPM NPNGRPNSDV LP.PWRNGMDV ARR...RDM
mmelre121P G.GAFDVSSE IAREWLRLPL NPNGRPNSDV LKPWRNAMDM TRRS...SDK
mmelre121P G.GAFDIPGD LARSWLSVP. NPDRVSNADV LKPWVNGMDL TRRP...SGR
mmelre121P GHKGFLLDGL EAGQMLSAE. ....RKNAEV IFPYLTGDEL LRTSPHPTR

901
mmelfeLORF3P WLVNVTP... .KQLR..SMP L..VLKRVE. QCKENR.LSG APDRQKLAAT 950
mmelre121P WLKDANP... .KDIH..QSP F..ILDRIN. KVAEFRSQOK SKDTQKYAKR
mmelre121P WLDG.ADIS. .EIR..AMP L..VLARIK. KVQEFRLKSS AKPTRQSAST
mmelre121P WFHGVSDVKR NHDLK..QMP Q..VQARIQ. AVKTMREASS DKQTQKDAAT
mmelre121P WLNE....AK YEKIK..SNP L..IQERIS. ISKNHREKST DKGTNKLALT
mmelre121P WLVDVAP... .SDIA..QSP V..LKKRLE. AVKSFRADSK AASTRKMAET
mmelre121P WISD....SQ EQEAK...SNS D..INCKLN. AVAAFRLLKSP KAATKKGAAW
mmelre121P WIND....DQ VDDAK..AIA E..IAKVLE. SCRSYR.QGA GRDAQKAANR
mmelre121P WIVDFGWEMS EQEAAALYAP FQHIREHVFP .....ER SKN.....
mmelre121P WIIDFGWTMS EADAALFETP FRHVLLHVFP .....ER DRN.....
mmelre121P WIIDFA.QMD EGEARQYLQP MAYVEQKIRP .....ER ATN.....
mmelre121P YVIDF.QPRD VFGARAYKLP FARIEREVLV TRQAAAAEEE ARNAEVLAAAN

```

mmelfeLORF3P	P.....	.....HL	.....FREQ	MNPDNYMIVP	LVTGCRRKYV
mmelre121P	P.....	.....ML	PTRLAYYSHD	EHTD.MLIVP	ATSSQRREYL
mmel	P.....	.....M.	..KFFYISQP	.DTDY.LLIP	ETSSENRQFI
mmelNMA1791	P.....	.....W.	..LFQKIRQP	SDGNY.LIIP	SVSSESRRFI
mmelBSU0677	P.....	.....W.	..KFRDTHE.	.TTNYSIVVP	SVSSENRFYI
mmelnegcry	P.....	.....HL	FGQ....RSQ	PDTDY.LCLP	KVVSERRSYF
mmelPflQ8	P.....	.....H.	..KFEEVKQI	GN.EVVTIVP	KVSSSESREYL
saro3834	P.....	.....HS	FC.YRTFQE.	...NIGIHVG	LTIGNGLSHV
mmelMSI135	....RRDAYR	...ERWWRHV	EPRPAFHASL	QGHSTRYMATP	RV.AKHRTFV
mmelCC0826	....NREMYR	...LNWVKHV	EPRQGLMKRV	PALSRLLVTP	EV.SKHRLFI
mmelDR0119.1	....SDRPSR	...ERWWLHQ	RSRPELREAT	IELDRFIGIP	RV.AKHLIPV
mmelDR2267	PKAKTNKHHR	NFLNQWWALS	YGRSEMIKI	SSLSTRYIVCS	RV.TKRQVFE

1001

mmelfeLORF3P	PFGYLG.NDI	IPTNLATIIP	EADHYAFGVL	ESIVHMAWMR	VVA...GRKG
mmelre121P	PIGYVSEKNI	VSYSL.MLIP	NASNPNFGIL	ESKVHYIWLK	NFC...GRLK
mmel	PIGFVDRNVI	SS.NATYHIP	SAEPLIFGLL	SSTMHNCWMR	NVG...GRLE
mmelNMA1791	PIGYLSFETV	VS.NLAFILP	NATLYHFGIL	SSTMHNAFMR	TVA...GRLK
mmelBSU0677	PMGLAGADTI	LS.NLIYVIY	DAEIYLLGIL	MSRMHMTWVK	AVA...GRLK
mmelnegcry	TVQRYPSNVI	AS.DLVFHAQ	DPDGLMFALA	SSSMFITWQK	SIG...GRLK
mmelPflQ8	PVGLLPRGSI	VT.DLAFALY	DAPLWNMALI	ASRLHLVWIG	~~~~~
saro3834	PAD.LKSSGF	VSSHTAYMIY	GWHPVEFALL	NSRLMLVWTE	TVG...GRLG
mmelMSI135	WLD....QAI	VPDSRIFAFS	RSDDVFFGIL	HSRFHEAWSF	GTCSWHGV.G
mmelCC0826	WLD....ARV	LPDHKLQVVT	LDDDCSFGVL	HSRFHEVWAL	AAGSWHGS.G
mmelDR0119.1	WLP....EGT	LPDSQVVVIA	RDDDFIFGVL	ASTIHRSWAR	MQGTYMGV.G
mmelDR2267	FLD....NGI	RPSDGLQIFA	FEDDYSFGVI	QSSVHWQWLI	ARG...GTLT

1051

mmelfeLORF3P	TSYRYSKNLV	YTNFPWPV.V	DINQKEK...	.....ITI	TAQDILNARN
mmelre121P	SDYRYSNTII	YNNFPWPT.V	GDKQEQN...	.....ISE	TAQGILNTRK
mmel	SRYRYSASLV	YNTFPWIQ..	....PNEKQ.	....SKAIEE	AAFAILKARS
mmelNMA1791	SDYRYSNTVV	YNNFPFPE..	SCRLPSENDER	PDPLRAAVEA	AAQTVLDARG
mmelBSU0677	TDYRYSAGLC	YNTFFPIPE.L	STRRKNE...	.....IEE	AILEILDLR.
mmelnegcry	SDLRFANTLT	WNTFPVPE.L	DEKTRQR...	.....IIK	AGKKVLDARA
mmelPflQ8	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
saro3834	NGMRFSNTIV	YNTFPVPS.L	TDQNK.....	....ADLTR	CAEDILLARE
mmelMSI135	NDPTYNSAGV	FETFPFPEGL	TPDIPAVRYE	KDSRAIAISK	AAKRLDDIRN
mmelCC0826	NDPRYTISTT	FETFPFPEGL	TPNIAAVDYE	GDPRQAIAA	AAAELNRLRE
mmelDR0119.1	NDLRYTPSTC	FETFPVP...	APT.....	.DEQRAEIEK	WAKYIVQLRE
mmelDR2267	ARLMYTSDTV	FDTFPWP*D.	.PTLAQVR..	....AVAA	AAVKLRELNR

1101

mmelfeLORF3P	.....	.....	...LY.....	....PDSSLA	DLYDPLTMPI
mmelre121P	.....	.....	...LY.....	....PDSSLA	DLYDPLTMPV
mmel	.....	.....	...NY.....	....PNESLA	GLYDPKTMPS
mmelNMA1791	.....	.....	...QYRREAQ	EAGLPEPTLA	ELYAPDAGYT
mmelBSU0677	.....	.....	.....EE	QGG....TLA	ELYNPSTMPI
mmelnegcry	.....	.....	...LH.....	....PERSLA	EHYNPLAMAP
mmelPflQ8	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
saro3834	.....	.....	...SHFP...	....ATIA	DLYDPETMPE
mmelMSI135	AWLNPSDLVQ	IKPEVVPGY	DRILPKDIAS	DAILRDRTLT	NLYNR.....
mmelCC0826	AWLNPPDLVR	IEPEVVPGY	DRVLPVSPEA	GAELKKRTLT	NLYNQ.....
mmelDR0119.1	HLLN.....	.....	.....Q	DA...KGTLT	GIYNQLEKLR
mmelDR2267	KVMREQ....	.....GW.	.....	.....SLR	DLYRTLDMGP

1151

mmelfeLORF3P	E.....	.LRKAHEAND	KAVLKAYGLK	P.SATEPEI.	.VQHLFKMYE
mmelre121P	E.....	.LRKAHEAND	KAVLKAYGLS	P.KATEQEI.	.VEHLFKMYE

1200

mme	E.....	.LLKAHQKLD	KAVDSVYGFK	.GPNT..EIA	RIAFLFETYQ
mmeNMA1791	A.....	.LDKAHATLD	KAVDKAYGYK	TGKNTDDEAE	RVAFLFELYR
mmeBSU0677	E.....	.LKVAHEKLD	GIVERAYRQK	QFES..DE.E	RLEVLLKLYQ
mmeegcry	E.....	.LIKAHDALD	REVDKAFGAP	RKLTTVRQ..	RQELLFANYE
mmePflQ8	-----	-----	-----	-----	-----
saro3834	S.....	.LRAAHRND	EVLERIY...	IGRRFRNDTE	RLEKLFELYT
mmeMSI135	.....RP.Q	WLVDASDLD	AAVAGAYGWP	ADIS...EDE	ALANLLELNL
mmeCC0826	.....RP.A	WLDMAHQRLD	AAVAAAYGWP	DGLT...DDE	ILERLFALNQ
mmeDR0119.1	NSPDAHPVS	ALATAHDKLD	QAVATAYGWE	WPLN...EDQ	VLERLLALNL
mmeDR2267	KNP.....	.LRDAQERLD	AAVSAAYGLP	AGA.....D	MLDFFLLALNA
	1201				1250
mmelfeLORF3P	KLTKKDW~~~	~~~~~	~~~~~	~~~~~	~~~~~
mmeLrel21P	KLTKGER*~~	~~~~~	~~~~~	~~~~~	~~~~~
mme	KMTSLLPPEK	EIKKSKGKN*	~~~~~	~~~~~	~~~~~
mmeNMA1791	KAAAI~	~~~~~	~~~~~	~~~~~	~~~~~
mmeBSU0677	EMTER~~~	~~~~~	~~~~~	~~~~~	~~~~~
mmeegcry	KLISHQP~~~	~~~~~	~~~~~	~~~~~	~~~~~
mmePflQ8	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
saro3834	KMTGGRSSEG	GAA~~~	~~~~~	~~~~~	~~~~~
mmeMSI135	AREAFNEHAK	SGLKTRKPRR	RPTPEEVRR	PQMKLPIAGG	RKSVVGPQQL
mmeCC0826	ERAAAGR~~~	~~~~~	~~~~~	~~~~~	~~~~~
mmeDR0119.1	ERCPA~~~	~~~~~	~~~~~	~~~~~	~~~~~
mmeDR2267	XVAAAEARGA	AVTGPGLPAG	LNTADFVTAD	AVRPLG*~~~	~~~~~
	1251		1273		
mmelfeLORF3P	~~~~~	~~~~~	~~~~~		
mmeLrel21P	~~~~~	~~~~~	~~~~~		
mme	~~~~~	~~~~~	~~~~~		
mmeNMA1791	~~~~~	~~~~~	~~~~~		
mmeBSU0677	~~~~~	~~~~~	~~~~~		
mmeegcry	~~~~~	~~~~~	~~~~~		
mmePflQ8	~~~~~	~~~~~	~~~~~		
saro3834	~~~~~	~~~~~	~~~~~		
mmeMSI135	TTKDRENQPT	SAERPRNTR	RTS		
mmeCC0826	~~~~~	~~~~~	~~~~~		
mmeDR0119.1	~~~~~	~~~~~	~~~~~		
mmeDR2267	~~~~~	~~~~~	~~~~~		